

# Guidelines on **Male Sexual Dysfunction:**

## **Erectile dysfunction and premature ejaculation**

E. Wespes (chair), I. Eardley, F. Giuliano, D. Hatzichristou,  
K. Hatzimouratidis (vice-chair), I. Moncada, A. Salonia, Y. Vardi

# TABLE OF CONTENTS

# PAGE

1.	BACKGROUND	4
1.1	Introduction	4
1.2	Methodology	4
1.3	Level of evidence and grade of recommendation	4
1.4	Publication History	5
1.5	Potential conflict of interest statement	5
1.6	References	5
2.	ERECTILE DYSFUNCTION	6
2.1	Epidemiology and risk factors	6
2.1.1	Epidemiology	6
2.1.2	Risk factors	6
2.1.3	Post-radical prostatectomy ED, post-radiotherapy ED & post-brachytherapy ED	6
2.1.4	Managing ED: implications for everyday clinical practice	7
2.1.5	Conclusions on the epidemiology of ED	7
2.1.6	References	7
2.2	Diagnostic evaluation	10
2.2.1	Basic work-up	10
2.2.1.1	Sexual history	11
2.2.1.2	Physical examination	11
2.2.1.3	Laboratory testing	11
2.2.2	Cardiovascular system and sexual activity: the patient at risk	12
2.2.2.1	Low-risk category	14
2.2.2.2	Intermediate- or indeterminate-risk category	14
2.2.2.3	High-risk category	14
2.2.3	Specialised diagnostic tests	14
2.2.3.1	Nocturnal penile tumescence and rigidity test	14
2.2.3.2	Intracavernous injection test	14
2.2.3.3	Duplex ultrasound of the penis	14
2.2.3.4	Arteriography and dynamic infusion cavernosometry or cavernosography	14
2.2.3.5	Psychiatric assessment	14
2.2.3.6	Penile abnormalities	14
2.2.4	Patient education - consultation and referrals	14
2.2.5	Guidelines for the diagnostic evaluation of ED	15
2.2.6	References	15
3.	TREATMENT OF ERECTILE DYSFUNCTION	17
3.1	Treatment options	17
3.2	Lifestyle management in ED with concomitant risk factors	18
3.3	Erectile dysfunction after radical prostatectomy	19
3.4	Causes of ED that can be potentially treated with a curative intent	20
3.4.1	Hormonal causes	20
3.4.2	Post-traumatic arteriogenic ED in young patients	20
3.4.3	Psychosexual counselling and therapy	20
3.5	First-line therapy	20
3.5.1	Oral pharmacotherapy	20
3.5.1.1	Sildenafil	20
3.5.1.2	Tadalafil	20
3.5.1.3	Vardenafil	21
3.5.1.4	Choice or preference between the different PDE5 inhibitors	21
3.5.1.5	On-demand or chronic use of PDE5 inhibitors	21
3.5.1.6	Safety issues for PDE5 inhibitors	22
3.5.1.6.1	Cardiovascular safety	22
3.5.1.6.2	Nitrates are contraindicated with PDE5 inhibitors	22
3.5.1.6.3	Antihypertensive drugs	23
3.5.1.6.4	$\alpha$ -Blocker interactions	23
3.5.1.6.5	Dosage adjustment	23
3.5.1.7	Management of non-responders to PDE5 inhibitors	23

	3.5.1.7.1 Check that the patient has been using a licensed medication	23
	3.6.1.7.2 Check that the medication has been properly prescribed and correctly used	23
	3.5.1.7.3 Possible manoeuvres in patients correctly using a PDE5 inhibitor	24
	3.5.2 Vacuum erection devices	24
	3.5.3 Shockwave therapy	25
3.6	Second-line therapy	25
	3.6.1 Intracavernous injections	25
	3.6.1.1 Alprostadil	25
	3.6.1.2 Combination therapy	26
	3.6.1.3 Intraurethral alprostadil	26
3.7	Third-line therapy (penile prostheses)	26
	3.7.1 Efficacy and satisfaction rates	26
	3.7.2 Complications	27
	3.7.3 Conclusions	27
3.8	Guidelines for the treatment of ED	28
3.9	References	28
4.	PREMATURE EJACULATION	37
	4.1 Introduction	37
	4.2 Definition of PE	37
	4.2.1 Overview	37
	4.2.2 Classifications	37
	4.3 Epidemiology	38
	4.3.1 Prevalence	38
	4.3.2 Pathophysiology and risk factors	38
	4.4 Impact on quality of life	38
	4.5 Diagnosis	39
	4.5.1 Intravaginal ejaculatory latency time (IELT)	39
	4.5.2 PE assessment questionnaires	39
	4.5.3 Physical examination and investigations	39
	4.6 Recommendations on the diagnosis of PE	40
	4.7 References	40
	4.8 Treatment	43
	4.8.1 Psychological/behavioural strategies	43
	4.8.1.1 Recommendation on the psychological/behavioural treatment of PE	44
	4.8.2 Topical anaesthetic agents	44
	4.8.2.1 Lidocaine-prilocaine cream	44
	4.8.2.2 SS-cream	44
	4.8.2.3 Recommendations on the topical therapy for PE	44
	4.8.3 Selective serotonin reuptake inhibitors	44
	4.8.3.1 Dapoxetine	45
	4.8.3.2 Recommendation on the treatment of PE	46
	4.8.4 Phosphodiesterase type 5 inhibitors	46
	4.8.4.1 Recommendation on the use of PDE5 inhibitors for the treatment of PE	46
	4.8.5 Other drugs	46
	4.8.6 Recommendations on the treatment of PE	46
	4.9 References	49
5.	CONCLUSION	52
6.	ABBREVIATIONS USED IN THE TEXT	53

# 1. BACKGROUND

## 1.1 Introduction

Erectile dysfunction (ED) and premature ejaculation (PE) are the two main complaints in male sexual medicine (1,2). New oral therapies have completely changed the diagnostic and therapeutic approach to ED and the Guidelines Office of the European Association of Urology (EAU) has appointed an Expert Panel to update previously published EAU guidelines for ED or impotence (3,4).

## 1.2 Methodology

The update is based on a systemic literature search performed by the panel members. The MedLine database was searched using the major Medical Subject Headings (MeSH) terms “erectile dysfunction”, “sexual dysfunction” “ejaculation”. MedLine search was supplemented by the term “premature ejaculation” in all search fields. All articles published between January 2009 (previous update) and January 2013 were considered for review. The Expert Panel has also identified critical problems and knowledge gaps, setting priorities for future clinical research.

## 1.3 Level of evidence and grade of recommendation

References in the text have been assessed according to their level of scientific evidence (LE), and guideline recommendations have been graded follow the listings in Tables 1 and 2, based on the Oxford Centre for Evidence-based Medicine Levels of Evidence (5). Grading aims to provide transparency between the underlying evidence and the recommendation given.

**Table 1: Level of evidence\***

Level	Type of evidence
1a	Evidence obtained from meta-analysis of randomised trials.
1b	Evidence obtained from at least one randomised trial.
2a	Evidence obtained from one well-designed controlled study without randomisation.
2b	Evidence obtained from at least one other type of well-designed quasi-experimental study.
3	Evidence obtained from well-designed non-experimental studies, such as comparative studies, correlation studies and case reports.
4	Evidence obtained from expert committee reports or opinions or clinical experience of respected authorities.

*\*Modified from (5).*

It should be noted that when recommendations are graded, the link between the LE and grade of recommendation (GR) is not directly linear. Availability of RCTs may not necessarily translate into a grade A recommendation where there are methodological limitations or disparity in published results.

Alternatively, absence of high level of evidence does not necessarily preclude a grade A recommendation, if there is overwhelming clinical experience and consensus. There may be exceptional situations where corroborating studies cannot be performed, perhaps for ethical or other reasons and in this case unequivocal recommendations are considered helpful. Whenever this occurs, it is indicated in the text as “upgraded based on panel consensus”. The quality of the underlying scientific evidence - although a very important factor - has to be balanced against benefits and burdens, values and preferences, and costs when a grade is assigned (4-6).

The EAU Guidelines Office does not perform structured cost assessments, nor can they address local/national preferences in a systematic fashion. But whenever these data are available, the expert panel will include the information.

**Table 2: Grade of recommendation\***

Grade	Nature of recommendations
A	Based on clinical studies of good quality and consistency that addressed the specific recommendations, including at least one randomised trial.
B	Based on well-conducted clinical studies, but without randomised clinical trials.
C	Made despite the absence of directly applicable clinical studies of good quality.

*\*Modified from (5).*

## 1.4 Publication History

The first European Association of Urology (EAU) Guidelines on Erectile Dysfunction were published in 2000 (6) with subsequent updates in 2001, 2002, 2004, 2005 and 2009. In particular the 2009 document presented a significant update of the previous publication with the inclusion of the topic "Premature Ejaculation" and the text was renamed to "EAU Guidelines on Male Sexual Dysfunction" (7). In 2011 the expert panel decided to develop separate guidelines addressing Penile Curvature, which resulted in a publication in 2012 (8). Several scientific summaries have been published in the EAU scientific journal, European Urology (9-12). Quick reference documents (pocket guidelines) are available presenting the main findings of both the Male Sexual Dysfunction Guidelines and the Penile Curvature Guidelines. These documents follow the updating cycle of the underlying large texts. All material can be viewed and downloaded for personal use at the EAU website. The EAU website also includes a selection of translations and republications produced by national urological associations: <http://www.uroweb.org/guidelines/online-guidelines/>.

The chapters on Erectile Dysfunction were peer-reviewed prior to publication.

## 1.5 Potential conflict of interest statement

The expert panel have submitted potential conflict of interest statements which can be viewed on the EAU website: <http://www.uroweb.org/guidelines/online-guidelines/>.

## 1.6 References

1. Lindau ST, Schumm LP, Laumann EO, et al. N Engl J Med. 2007 Aug 23;357(8):762-74.  
<http://www.ncbi.nlm.nih.gov/pubmed/17715410>
2. Rosenberg MT, Sadovsky R. Identification and diagnosis of premature ejaculation. Int J Clin Pract 2007 Jun;61(6):903-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/17504352>
3. Wespes E, Amar E, Hatzichristou DG, et al. European Association of Urology Guidelines on erectile dysfunction. Eur Urol 2002 Jan;41(1):1-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/11999460>
4. Hatzimouratidis K, Amar E, Eardley I, et al; European Association of Urology. Guidelines on male sexual dysfunction: erectile dysfunction and premature ejaculation. Eur Urol. 2010 May;57(5):804-14.  
<http://www.ncbi.nlm.nih.gov/pubmed/20189712>
5. Oxford Centre for Evidence-based Medicine Levels of Evidence (May 2001). Produced by Bob Phillips, Chris Ball, Dave Sackett, Doug Badenoch, Sharon Straus, Brian Haynes, Martin Dawes since November 1998.  
<http://www.cebm.net/index.aspx?o=1025> [access date January 2013].
6. Wespes E, Amar E, Hatzichristou D, et al; EAU Guidelines Panel on Erectile Dysfunction. EAU Guidelines on Erectile Dysfunction. Edition presented at the EAU Annual Congress Brussels, 2000.
7. Wespes E, Amar E, Eardley I, et al; EAU Guidelines Panel on Male Sexual Dysfunction. EAU Guidelines on Male Sexual Dysfunction (Erectile Dysfunction and premature ejaculation). Edition presented at the EAU Annual Congress Stockholm, 2009. ISBN 978-90-79754-09-0.
8. Wespes E, Hatzimouratidis K, Eardley I, et al; EAU Guidelines Panel on Male Sexual Dysfunction. EAU Guidelines on Penile Curvature. Edition presented at the EAU Annual Congress Paris, 2012. ISBN 978-90-79754-83-0.
9. Wespes E, Amar E, Hatzichristou D, et al. Guidelines on Erectile Dysfunction. Eur Urol 2002 Jan 41(1):1-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/11999460>
10. Wespes E, Amar E, Hatzichristou D, et al; EAU. EAU Guidelines on erectile dysfunction: an update. Eur Urol. 2006 May;49(5):806-15.  
<http://www.ncbi.nlm.nih.gov/pubmed/16530932>
11. Hatzimouratidis K, Amar E, Eardley I, et al European Association of Urology. Guidelines on male sexual dysfunction: erectile dysfunction and premature ejaculation. Eur Urol. 2010 May;57(5):804-14.  
<http://www.ncbi.nlm.nih.gov/pubmed/20189712>
12. Hatzimouratidis K, Eardley I, Giuliano F, et al; European Association of Urology. EAU guidelines on penile curvature. Eur Urol. 2012 Sep;62(3):543-52.  
<http://www.ncbi.nlm.nih.gov/pubmed/22658761>

## 2. ERECTILE DYSFUNCTION

### 2.1 Epidemiology and risk factors

Erection is a neuro-vasculo-tissular phenomenon under hormonal control. It includes arterial dilatation, trabecular smooth muscle relaxation, and activation of the corporeal veno-occlusive mechanism (1,2).

Erectile dysfunction is defined as the persistent inability to attain and maintain an erection sufficient to permit satisfactory sexual performance. Although ED is a benign disorder, it may affect physical and psychosocial health and may have a significant impact on the quality of life (QoL) of sufferers and their partners (3). There is increasing evidence that ED can be an early manifestation of coronary artery and peripheral vascular disease; thus, ED should not be regarded only as a QoL issue but also as a potential warning sign of cardiovascular disease (4-8).

#### 2.1.1 Epidemiology

Epidemiological data have shown a high prevalence and incidence of ED worldwide. The first large, community-based study of ED was the Massachusetts Male Aging Study (MMAS) (3). The study reported an overall prevalence of 52% ED in non-institutionalised 40-70-year-old men in the Boston area; specific prevalence for minimal, moderate, and complete ED was 17.2%, 25.2%, and 9.6%, respectively. In the Cologne study of men aged 30-80 years, the prevalence of ED was 19.2%, with a steep age-related increase from 2.3% to 53.4% (9). In the National Health and Social Life Survey (NHSLs), the prevalence of sexual dysfunction in males (not specific ED) was 31% (10). The incidence rate of ED (new cases per 1,000 men annually) was 26 in the MMAS study (11), 65.6 (mean follow-up of 2 years) in a Brazilian study (12), and 19.2 (mean follow-up of 4.2 years) in a Dutch study (13). In Taiwan, the prevalence of ED was 27% among all patients investigated and 29% among those aged  $\geq 40$  years (14). In Ghana, the overall prevalence of ED is 59.6% and there are positive correlations between ED, dissatisfaction, age and other sexual dysfunctions (15). Differences between these studies can be explained by differences in methodology and in the ages, socioeconomic and cultural status of the populations studied.

Data from epidemiological studies have demonstrated consistent and compelling evidence for an association between lower urinary tract symptoms (LUTS)/benign prostatic hypertrophy (BPH) and sexual dysfunction in aging men that is independent of the effects of age, other comorbidities, and various lifestyle factors (16). The Massachusetts Male Aging (MSAM-7) study systematically investigated the relationship between LUTS and sexual dysfunction in  $> 12,000$  men aged 50-80 years. It was performed in the US and six European countries (France, Germany, Italy, Netherlands, Spain, and UK). Eighty-three percent of men considered themselves sexually active, and 71% reported at least one episode of sexual activity in the past 4 weeks. The overall prevalence of LUTS was 90%. Only 19% of men had sought medical help for LUTS and only 11% were medically treated. The overall prevalence of ED was 49%, and 10% of patients reported complete absence of erection. The overall prevalence of ejaculation disorders was 46% and 5% reported anejaculation (17).

#### 2.1.2 Risk factors

Erectile dysfunction shares common risk factors with cardiovascular disease (e.g., lack of exercise, obesity, smoking, hypercholesterolaemia, and metabolic syndrome); some of which can be modified. Moreover, men with mild ED have similar risk factors to those of a general ED clinical trial population. Thus, mild ED is an important indicator of risk for associated underlying disease. Men complaining of mild ED should be evaluated adequately (for underlying cardiovascular risk) (18).

In the MMAS, men who began exercising in midlife had a 70% reduced risk for ED compared to sedentary men and a significantly lower incidence over an 8-year follow-up period of regular exercise (19). A multicentre, randomised, open-label study in obese men with moderate ED compared 2 years of intensive exercise and weight loss with a control group given general information about healthy food choices and exercise (20). Significant improvements in body mass index (BMI) and physical activity scores, as well as erectile function, were observed in the lifestyle intervention group. These changes were highly correlated with both weight loss and activity levels.

Some studies have shown some evidence that lifestyle modification and pharmacotherapy for cardiovascular risk factors are effective in improving sexual function in men with ED. However, it should be emphasized that more controlled prospective studies are necessary to determine the effects of exercise or other lifestyle changes in prevention or treatment of ED (6).

#### 2.1.3 Post-radical prostatectomy ED, post-radiotherapy ED & post-brachytherapy ED

Radical prostatectomy (RP) in any form (open, laparoscopic, or robotic) is a widely performed procedure for patients with clinically localised prostate cancer and a life expectancy of at least 10 years. This procedure may lead to treatment-specific sequelae affecting health-related QoL. This outcome has become increasingly

important with the more frequent diagnosis of prostate cancer in younger patients (21-22). Research has shown that 25-75% of men experience postoperative ED (23). A systematic review has shown that incidence of potency recovery after robotic prostatectomy is influenced by numerous factors. It reported, for the first time, a significant advantage in favor of robotic laparoscopic radical prostatectomy in comparison with retropubic radical prostatectomy in terms of 12-month potency rates (24). However, there was no significant difference between laparoscopic RP and robot-assisted laparoscopic RP. Today, we do not have enough evidence-based data to consider that robot-assisted laparoscopic RP has any advantageous effect on functional outcome. Experience of the surgeon seems to be the main factor besides preservation of neurovascular bundles and patient age.

Post-RP ED is multifactorial. Cavernal nerve injury induces proapoptotic (loss of smooth muscle) and profibrotic (increase in collagen) factors within the corpora cavernosa. These changes may also be caused by poor oxygenation due to changes in the blood supply to the cavernosa because of possible arterial damage during the surgical procedure.

Preoperative potency is a major factor associated with the recovery of erectile function after surgery, therefore, patients being considered for nerve-sparing radical prostatectomy (NSRP) should ideally be potent preoperatively (24-29). It is also clear that cavernosal nerves must be preserved to ensure erectile function recovers after RP. In addition, the role of vascular insufficiency is of increasing interest in postoperative ED (30,31).

ED is also a common sequela after external beam radiotherapy and brachytherapy for prostate cancer. The mechanisms contributing to ED after prostate irradiation involve injury to the neurovascular bundles, penile vasculature, and cavernosal structural tissue (32,33). Alternative treatments for prostate cancer including cryotherapy and high-intensity focused ultrasound are associated with equivalent or worsened rates of ED compared to surgery or radiation therapy (34,35).

#### 2.1.4 **Managing ED: implications for everyday clinical practice**

Advances in basic and clinical research in ED during the past 15 years have led to the development of a variety of new treatment options, including pharmacological agents for intracavernous, intraurethral, and oral use (36-38). Reconstructive vascular surgery is reserved for select cases of arterial insufficiency, with no current indications for venous ligation procedures, given the poor overall outcomes (39,40).

An increasing number of men are currently seeking help for ED due to the growing public awareness of the condition and the availability of effective, safe and user-friendly oral drug therapy. However, not all physicians evaluating and treating ED have appropriate background knowledge and clinical experience in sexual medicine. Thus, men with ED may receive little or no evaluation before treatment and will therefore not receive treatment for any underlying disease that may be causing their ED. Other men without ED may be requesting treatment simply to enhance their sexual performance.

#### 2.1.5 **Conclusions on the epidemiology of ED**

	<b>LE</b>
Erection is a neuro-vasculo-tissular phenomenon under hormonal control.	2b
ED is common worldwide.	2b
ED shares risk factors with cardiovascular disease.	2b
Lifestyle modification (intensive exercise and decrease in BMI) can improve erectile function.	1b
ED is a symptom, not a disease. Some patients may not be properly evaluated or receive treatment for an underlying disease or condition that may be causing ED.	4
ED is common after radical prostatectomy, irrespective of the surgical technique used.	2b
ED is common after external radiotherapy and brachytherapy.	2b

#### 2.1.6 **References**

1. Lue TF, Tanagho EA. Physiology of erection and pharmacological management of impotence. J Urol 1987 May;137(5):829-36.  
<http://www.ncbi.nlm.nih.gov/pubmed/3553617>
2. Gratzke C, Angulo J, Chitale K, Dai YT, Kim NN, Paick JS, Simonsen U, Uckert S, Wespes E, Andersson KE, Lue TF, Stief CG. Anatomy, physiology, and pathophysiology of erectile dysfunction. J Sex Med. 2010 Jan;7(1 Pt 2):445-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/20092448>
3. Feldman HA, Goldstein I, Hatzichristou DG, et al. Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Aging Study. J Urol 1994 Jan;151(1):54-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/8254833>

4. Jackson G, Boon N, Eardley I, et al. Erectile dysfunction and coronary artery disease prediction: evidence-based guidance and consensus. *Int J Clin Pract.* 2010 Jun;64(7):848-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/20584218>
5. Dong JY, Zhang YH, Qin LQ. Erectile dysfunction and risk of cardiovascular disease: Meta-analysis of prospective cohort studies. *J Am Coll Cardiol* 2011 Sep 20;58(13):1378-85.  
<http://www.ncbi.nlm.nih.gov/pubmed/21920268>
6. Gupta BP, Murad MH, Clifton MM, et al. The effect of lifestyle modification and cardiovascular risk factor reduction on erectile dysfunction: a systematic review and meta-analysis. *Arch Intern Med.* 2011 Nov 14;171(20):1797-803.  
<http://www.ncbi.nlm.nih.gov/pubmed/21911624>
7. Guo W, Liao C, Zou Y, Li F, Li T, Zhou Q, Cao Y, Mao X. Erectile dysfunction and risk of clinical cardiovascular events: A meta-analysis of seven cohort studies. *J Sex Med* 2010 Aug;7(8):2805-16.  
<http://www.ncbi.nlm.nih.gov/pubmed/20367771>
8. Batty GD, Li Q, Czernichow S, Neal B, et al. Erectile dysfunction and later cardiovascular disease in men with type 2 diabetes: Prospective cohort study based on the ADVANCE (Action in Diabetes and Vascular Disease: Preterax and Diamicron Modified-Release Controlled Evaluation) trial. *J Am Coll Cardiol* 2010 Nov 30;56(23):1908-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/21109113>
9. Braun M, Wassmer G, Klotz T, et al. Epidemiology of erectile dysfunction: results of the 'Cologne Male Survey'. *Int J Impot Res* 2000 Dec;12(6):305-11.  
<http://www.ncbi.nlm.nih.gov/pubmed/11416833>
10. Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999 Feb;281(6):537-44.  
<http://www.ncbi.nlm.nih.gov/pubmed/10022110>
11. Johannes CB, Araujo AB, Feldman HA, et al. Incidence of erectile dysfunction in men 40 to 69 years old: longitudinal results from the Massachusetts Male Aging Study. *J Urol* 2000 Feb;163(2):460-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/10647654>
12. Moreira ED Jr, Lobo CF, Diamant A, et al. Incidence of erectile dysfunction in men 40 to 69 years old: results from a population-based cohort study in Brazil. *Urology* 2003 Feb;61(2):431-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12597962>
13. Schouten BW, Bosch JL, Bernsen RM, et al. Incidence rates of erectile dysfunction in the Dutch general population. Effects of definition, clinical relevance and duration of follow-up in the Krimpen Study. *Int J Impot Res* 2005 Jan-Feb;17(1):58-62.  
<http://www.ncbi.nlm.nih.gov/pubmed/15510192>
14. Hwang TI, Tsai TF, Lin YC, et al. A survey of erectile dysfunction in Taiwan: use of the erection hardness score and quality of erection questionnaire. *J Sex Med* 2010 Aug;7(8):2817-24.  
<http://www.ncbi.nlm.nih.gov/pubmed/20456624>
15. Amidu N, Owiredu WK, Woode E, et al. Prevalence of male sexual dysfunction among Ghanaian populace: myth or reality? *Int J Impot Res* 2010 Nov-Dec;22(6):337-42.  
<http://www.ncbi.nlm.nih.gov/pubmed/20927122>
16. Seftel AD, de la Rosette J, Birt J, et al. Coexisting lower urinary tract symptoms and erectile dysfunction: a systematic review of epidemiological data. *Int J Clin Pract* 2013 Jan;67(1):32-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/23082930>
17. Rosen R, Altwein J, Boyle P, et al. Lower urinary tract symptoms and male sexual dysfunction: the multinational survey of the aging male (MSAM-7). *Eur Urol* 2003 Dec;44(6):637-49.  
<http://www.ncbi.nlm.nih.gov/pubmed/14644114>
18. Lee JC, Bénard F, Carrier S, et al. Do men with mild erectile dysfunction have the same risk factors as the general erectile dysfunction clinical trial population? *BJU Int* 2011 Mar;107(6):956-60.  
<http://www.ncbi.nlm.nih.gov/pubmed/20950304>
19. Derby CA, Mohr BA, Goldstein I, et al. Modifiable risk factors and erectile dysfunction: can lifestyle changes modify risk? *Urology* 2000 Aug;56(2):302-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/10925098>
20. Esposito K, Giugliano F, Di Palo C, et al. Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled trial. *JAMA* 2004 Jun;291(24):2978-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/15213209>
21. Salonia A, Burnett AL, Graefen M, et al. Prevention and management of postprostatectomy sexual dysfunctions. Part 1: choosing the right patient at the right time for the right surgery. *Eur Urol.* 2012 Aug;62(2):261-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/22575909>



22. Salonia A, Burnett AL, Graefen M, et al. Prevention and management of postprostatectomy sexual dysfunctions part 2: recovery and preservation of erectile function, sexual desire, and orgasmic function. *Eur Urol* 2012 Aug;62(2):273-86.  
<http://www.ncbi.nlm.nih.gov/pubmed/22575910>
23. Sanda MG, Dunn RL, Michalski J, et al. Quality of life and satisfaction with outcome among prostatecancer survivors. *N Engl J Med* 2008 Mar;358(12):1250-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/18354103>
24. Ficarra V, Novara G, Ahlering TE, et al. Systematic review and meta-analysis of studies reporting potency rates after robot-assisted radical prostatectomy. *Eur Urol*. 2012 Sep;62(3):418-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/22749850>
25. Hatzimouratidis K, Burnett AL, Hatzichristou D, et al. Phosphodiesterase type 5 inhibitors in postprostatectomy erectile dysfunction: a critical analysis of the basic science rationale and clinical application. *Eur Urol* 2009, 55:334-347.  
<http://www.ncbi.nlm.nih.gov/pubmed/18986755>
26. Magheli A, Burnett AL. Erectile dysfunction following prostatectomy: prevention and treatment. *Nat Rev Urol* 2009 Aug;6(8):415-27.  
<http://www.ncbi.nlm.nih.gov/pubmed/19657376>
27. Ferronha F, Barros F, Vaz Santos V, et al. Is there any evidence of superiority between retropubic, laparoscopic or robot-assisted radical prostatectomy? *International Braz J Urol* 2011 March-April;37(2):146-60.  
<http://www.ncbi.nlm.nih.gov/pubmed/21557832>
28. Barry MJ, Gallagher PM, Skinner JS, et al. Adverse effects of robotic-assisted laparoscopic versus open retropubic radical prostatectomy among a nationwide random sample of medicare-age men. *J Clin Oncol* 2012 Feb 10;30(5):513-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/22215756>
29. Vickers A, Savage C, Bianco F, et al. Cancer control and functional outcomes after radical prostatectomy as markers of surgical quality: analysis of heterogeneity between surgeons at a single cancer center. *Eur Urol* 2011 Mar;59(3):317-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/21095055>
30. Mulhall JP, Slovick R, Hotaling J, et al. Erectile dysfunction after radical prostatectomy: hemodynamic profiles and their correlation with the recovery of erectile function. *J Urol* 2002 Mar;167(3):1371-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/11832735>
31. Secin FP, Touijer K, Mulhall J, et al. Anatomy and preservation of accessory pudendal arteries in laparoscopic radical prostatectomy. *Eur Urol* 2007 May;51(5):1229-35.  
<http://www.ncbi.nlm.nih.gov/pubmed/16989942>
32. van der Wielen GJ, Mulhall JP, Incrocci L. Erectile dysfunction after radiotherapy for prostate cancer and radiation dose to the penile structures: a critical review. *Radiother Oncol* 2007 Aug;84(2):107-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/17707936>
33. Stember DS, Mulhall JP. The concept of erectile function preservation (penile rehabilitation) in the patient after brachytherapy for prostate cancer. *Brachytherapy* 2012 Mar-Apr;11(2):87-96.  
<http://www.ncbi.nlm.nih.gov/pubmed/22330103>
34. Cordeiro ER, Cathelineau X, Thuroff S, et al. High-intensity focused ultrasound (HIFU) for definitive treatment of prostate cancer. *BJU Int*. 2012 Nov;110(9):1228-42.  
<http://www.ncbi.nlm.nih.gov/pubmed/22672199>
35. Williams SB, Lei Y, Nguyen PL, et al. Comparative effectiveness of cryotherapy vs brachytherapy for localised prostate cancer. *BJU Int*. 2012 Jul;110(2 Pt 2):E92-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/22192688>
36. Goldstein I, Lue TF, Padma-Nathan H, et al; Sildenafil Study Group. Oral sildenafil in the treatment of erectile dysfunction. 1998. *J Urol* 2002 Feb;167(2 Pt 2):1197-203.  
<http://www.ncbi.nlm.nih.gov/pubmed/11905901>
37. Hellstrom WJ, Gittelman M, Karlin G, et al; Vardenafil Study Group. Sustained efficacy and tolerability of vardenafil, a highly potent selective phosphodiesterase type 5 inhibitor, in men with erectile dysfunction: results of a randomized, double-blind, 26-week placebo-controlled pivotal trial. *Urology* 2003 Apr;61(4 Suppl 1):8-14.  
<http://www.ncbi.nlm.nih.gov/pubmed/12657355>
38. Brock GB, McMahon CG, Chen KK, et al. Efficacy and safety of tadalafil for the treatment of erectile dysfunction: results of integrated analyses. *J Urol* 2002 Oct;168(4 Pt 1):1332-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12352386>

39. Wespes E, Schulman C. Venous impotence: pathophysiology, diagnosis and treatment. J Urol 1993 May;149(5 Pt 2):1238-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/8479008>
40. Rao DS, Donatucci CF. Vasculogenic impotence. Arterial and venous surgery. Urol Clin North Am 2001 May;28(2):309-19.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402583>

## 2.2 Diagnostic evaluation

### 2.2.1 Basic work-up

The first step in evaluating ED is always a detailed medical and sexual history of patients and partners when available (1,2). Often it is not possible to include the partner on the patient's first visit, but an effort should be made to include the partner at the second visit. The pathophysiology of ED may be vasculogenic, neurogenic, anatomical, hormonal, drug-induced and/or psychogenic (Table 3) (3). Taking a comprehensive medical history may reveal one of the many common disorders associated with ED.

It is important to establish a relaxed atmosphere during history-taking. This will make it easier to ask questions about erectile function and other aspects of sexual history. A relaxed atmosphere will also make it easier to explain the diagnosis and therapeutic approach to the patient and his partner.

**Table 3: Pathophysiology of ED**

<b>Vasculogenic</b>	
-	Cardiovascular disease
-	Hypertension
-	Diabetes mellitus
-	Hyperlipidaemia
-	Smoking
-	Major surgery (RP) or radiotherapy (pelvis or retroperitoneum)
<b>Neurogenic</b>	
<i>Central causes</i>	
-	Degenerative disorders (multiple sclerosis, Parkinson's disease, multiple atrophy etc.)
-	Spinal cord trauma or diseases
-	Stroke
-	Central nervous system tumours
<i>Peripheral causes</i>	
-	Type 1 and 2 diabetes mellitus
-	Chronic renal failure
-	Polyneuropathy
-	Surgery (pelvis or retroperitoneum, radical prostatectomy, colorectal surgery, etc.)
<b>Anatomical or structural</b>	
-	Hypospadias, epispadias
-	Micropenis
-	Congenital curvature of the penis
-	La Peyronie's disease
<b>Hormonal</b>	
-	Hypogonadism
-	Hyperprolactinemia
-	Hyper- and hypothyroidism
-	Hyper- and hypocortisolism (Cushing's disease etc)
<b>Drug-induced</b>	
-	Antihypertensives (diuretics are the most common medication causing ED)
-	Antidepressants (selective serotonin reuptake inhibitors, tricyclics)
-	Antipsychotics (incl. neuroleptics)
-	Antiandrogens; GnRH analogues and antagonists
-	Recreational drugs (alcohol, heroin, cocaine, marijuana, methadone)
<b>Psychogenic</b>	
-	Generalised type (e.g., lack of arousability and disorders of sexual intimacy)
-	Situational type (e.g., partner-related, performance-related issues or due to distress)

<b>Trauma</b>	
-	Penile fracture

### 2.2.1.1 Sexual history

The sexual history must include (when available) information about previous and current sexual relationships, current emotional status, onset and duration of the erectile problem, and previous consultations and treatments. The sexual health status of the partner(s) (when available) can also be useful. A detailed description should be made of the rigidity and duration of both sexually stimulated and morning erections and of problems with arousal, ejaculation, and orgasm. Validated psychometric questionnaires, such as the International Index for Erectile Function (IIEF) (4), help to assess the different sexual function domains (i.e, sexual desire, erectile function, orgasmic function, ejaculation, intercourse, and overall satisfaction), as well as the impact of a specific treatment modality. Psychometric analysis also supports the use of erectile hardness score as a simple, reliable and valid tool for the assessment of penile rigidity in practice and in clinical trials research (5). In cases of clinical depression, the use of a 2-question scale for depression is recommended: "During the past month have you often been bothered by feeling down, depressed or hopeless? During the past month have you often been bothered by little interest or pleasure, doing things?" (6). Patients should be screened for symptoms of possible of possible hypogonadism, including decreased energy, libido, fatigue, and cognitive impairment, as well as for symptomatic lower urinary tract symptoms. Where indicated, screening questionnaires, such as the International Prostate Symptom Score may be utilised.

### 2.2.1.2 Physical examination

Every patient must be given a physical examination focused on the genitourinary, endocrine, vascular, and neurological systems (1). A physical examination may reveal unsuspected diagnoses, such as La Peyronie's disease, prostatic enlargement or irregularity/nodularity, or signs and symptoms suggesting hypogonadism (small testes, alterations in secondary sexual characteristics etc.) (2). A rectal examination should be performed in every patient older than 40 years. Blood pressure and heart rate should be measured if they have not been assessed in the previous 3-6 months. Particular attention must be given to patients with cardiovascular disease (Section 2.2.2).

### 2.2.1.3 Laboratory testing


Laboratory testing must be tailored to the patient's complaints and risk factors. Patients may need a fasting glucose or HbA1c and lipid profile if not recently assessed. Hormonal tests include a morning sample of total testosterone. If indicated bioavailable or calculated-free testosterone may be needed to corroborate total testosterone measurements. However, the threshold of testosterone to maintain ED is low and ED is usually a symptom of more severe cases of hypogonadism (7). For levels > 8 nmol/l the relationship between circulating testosterone and sexual function is very low (7,8).

Additional laboratory tests may be considered in selected patients, for example, prostate-specific antigen (PSA) for detection, or suspicion, of prostate cancer (9). Additional hormonal tests, for example, prolactin, and luteinizing hormone, are performed when low testosterone levels are detected. If any abnormality is observed, referral to an endocrinologist may be indicated (10,11).

Although physical examination and laboratory evaluation of most men with ED may not reveal the exact diagnosis, these opportunities to identify critical comorbid conditions should not be missed (12).

Figure 1 gives the minimal diagnostic evaluation (basic work-up) in patients with ED.

**Figure 1: Minimal diagnostic evaluation (basic work-up) in patients with ED**



ED = erectile dysfunction; IIEF = International Index of Erectile Function.

### 2.2.2 Cardiovascular system and sexual activity: the patient at risk

Patients who seek treatment for sexual dysfunction have a high prevalence of cardiovascular disease. The cardiac risks associated with sexual activity are well established. Epidemiological surveys have emphasised the association between cardiovascular and metabolic risk factors and sexual dysfunction in men and women (13). ED can improve the sensitivity of screening for asymptomatic cardiovascular disease in men with diabetes (14,15). ED significantly increases the risk of cardiovascular disease, coronary heart disease, stroke, and all-cause mortality, and the increase is probably independent of conventional cardiovascular risk factors (16).

There has been an extensive investigation of the pharmacological properties of phosphodiesterase 5 inhibitors (PDE5Is), including their effects on cardiac smooth muscle activity and overall cardiovascular safety. The EAU Guidelines recommendations given here for treating men with ED have been adapted from previously published recommendations from the Princeton Consensus conferences on sexual dysfunction and cardiac risk (17-19). The Princeton Consensus (Expert Panel) Conference is dedicated to optimizing sexual function and preserving cardiovascular health. A total of three consensus papers have been published (17-19). The Third Princeton Consensus had two primary objectives. The first focused on evaluation and management of cardiovascular risk in men with ED and no known cardiovascular disease, with particular emphasis on identification of men with ED who may require additional cardiological work-up. The second objective focused on re-evaluation and modification of previous recommendations for evaluation of cardiac risk associated with sexual activity in men with known cardiovascular disease. The recommendations built on those developed during the first and second Princeton Consensus Conferences; first, emphasising the use of exercise ability and stress testing to ensure that each man's cardiovascular health is consistent with the physical demands of sexual activity before prescribing treatment for ED; and second, highlighting the link between ED and


cardiovascular disease, which may be asymptomatic and benefit from cardiovascular risk reduction (19). Patients with ED can be stratified into three cardiovascular risk categories (Table 4), which can be used as the basis for a treatment algorithm for initiating or resuming sexual activity (Figure 2). It is also possible for the clinician to estimate the risk of sexual activity in most patients from their level of exercise tolerance, which can be determined when taking the patient's history.

**Table 4: Cardiac risk stratification (based on 2<sup>nd</sup> Princeton Consensus) (18)**

Low-risk category	Intermediate-risk category	High-risk category
Asymptomatic, < 3 risk factors for CAD (excluding sex)	≥ 3 risk factors for CAD (excluding sex)	High-risk arrhythmias
Mild, stable angina (evaluated and/or being treated)	Moderate, stable angina	Unstable or refractory angina
Uncomplicated previous MI	Recent MI (> 2, < 6 weeks)	Recent MI (< 2 weeks)
LVD/CHF (NYHA class I)	LVD/CHF (NYHA class II)	LVD/CHF (NYHA class III/IV)
Post-successful coronary revascularisation	Non-cardiac sequelae of atherosclerotic disease (e.g., stroke, peripheral vascular disease)	Hypertrophic obstructive and other cardiomyopathies
Controlled hypertension		Uncontrolled hypertension
Mild valvular disease		Moderate-to-severe valvular disease

CAD = coronary artery disease; CHF = congestive heart failure; LVD = left ventricular dysfunction; MI = myocardial infarction; NYHA = New York Heart Association.

**Figure 2: Treatment algorithm for determining level of sexual activity according to cardiac risk in ED (based on 3<sup>rd</sup> Princeton Consensus) (19)**



<sup>a</sup>Sexual activity is equivalent to walking 1 mile on the flat in 20 min or briskly climbing two flights of stairs in 10 s.

<sup>b</sup>Sexual activity is equivalent to 4 min of the Bruce treadmill protocol.

### 2.2.2.1 *Low-risk category*

The low-risk category includes patients who do not have any significant cardiac risk associated with sexual activity. Low risk is typically implied by the ability to perform exercise of modest intensity, which is defined as  $\geq 6$  "metabolic equivalents of energy expenditure in the resting state" (METs) without symptoms. According to current knowledge of the exercise demand or emotional stress associated with sexual activity, low-risk patients do not need cardiac testing or evaluation before the initiation or resumption of sexual activity or therapy for sexual dysfunction.

### 2.2.2.2 *Intermediate- or indeterminate-risk category*

The intermediate- or indeterminate-risk category consists of patients with an uncertain cardiac condition or patients whose risk profile requires testing or evaluation before the resumption of sexual activity. Based upon the results of testing, these patients may be moved to either the high- or low-risk group. A cardiology consultation may be needed in some patients to help the primary physician determine the safety of sexual activity.

### 2.2.2.3 *High-risk category*

High-risk patients have a cardiac condition that is sufficiently severe and/or unstable for sexual activity to carry a significant risk. Most high-risk patients have moderate-to-severe symptomatic heart disease. High-risk individuals should be referred for cardiac assessment and treatment. Sexual activity should be stopped until the patient's cardiac condition has been stabilised by treatment, or a decision made by the cardiologist and/or internist that it is safe to resume sexual activity.

## 2.2.3 **Specialised diagnostic tests**

Most patients with ED can be managed within the sexual care setting, conversely, some patients may need specific diagnostic tests (Tables 5 and 6).

### 2.2.3.1 *Nocturnal penile tumescence and rigidity test*

The nocturnal penile tumescence and rigidity (NPTR) assessment should be done on at least two nights. A functional erectile mechanism is indicated by an erectile event of at least 60% rigidity recorded on the tip of the penis that lasts for  $\geq 10$  min (20).

### 2.2.3.2 *Intracavernous injection test*

The intracavernous injection test gives limited information about vascular status. A positive test is a rigid erectile response (unable to bend the penis) that appears within 10 min after the intracavernous injection and lasts for 30 min (21). This response indicates a functional, but not necessarily normal, erection, and the erection may coexist with arterial insufficiency and/or veno-occlusive dysfunction (22). A positive test shows that a patient will respond to the intracavernous injection programme. The test is inconclusive as a diagnostic procedure and duplex Doppler study of the penis should be requested, if clinically warranted.

### 2.2.3.3 *Duplex ultrasound of the penis*

A peak systolic blood flow  $> 30$  cm/s, an end-diastolic velocity of  $< 3$  cm/s and a resistance index  $> 0.8$  are generally considered normal (21). Further vascular investigation is unnecessary when a Duplex examination is normal.

### 2.2.3.4 *Arteriography and dynamic infusion cavernosometry or cavernosography*

Arteriography and dynamic infusion cavernosometry or cavernosography (DICC) should be performed only in patients who are being considered for vascular reconstructive surgery (23).

### 2.2.3.5 *Psychiatric assessment*

Patients with psychiatric disorders must be referred to a psychiatrist who is particularly interested in ED. In younger patients ( $< 40$  years) with long-term primary ED, psychiatric assessment may be helpful before any organic assessment is carried out.

### 2.2.3.6 *Penile abnormalities*

Surgical correction may be needed for patients with ED due to penile abnormalities, e.g. hypospadias, congenital curvature, or Peyronie's disease with preserved rigidity.

## 2.2.4 **Patient education - consultation and referrals**

Consultation with the patient should include a discussion of the expectations and needs of both the patient and his stable sexual partner, if available. It should also review both the patient's and partner's understanding

of ED, the results of diagnostic tests, and provide a rational selection of treatment options. Patient and partner education is an essential part of ED management (24,25).

**Table 5: Indications for specific diagnostic tests**

Primary ED (not caused by organic disease or psychogenic disorder).
Young patients with a history of pelvic or perineal trauma who could benefit from potentially curative vascular surgery.
Patients with penile deformities that might require surgical correction, e.g., Peyronie's disease, congenital curvature.
Patients with complex psychiatric or psychosexual disorders.
Patients with complex endocrine disorders.
Specific tests may be indicated at the request of the patient or his partner.
Medicolegal reasons, e.g., implantation of penile prosthesis, sexual abuse.

**Table 6: Specific diagnostic tests**

NTPR using Rigiscan
Vascular studies
- Intracavernous vasoactive drug injection
- Duplex Doppler study of the penis
- Dynamic Infusion Caverosometry and Caverosography (DICC)
- Internal pudendal arteriography
Neurological studies, e.g., bulbocavernosus reflex latency, nerve conduction studies
Endocrinological studies
Specialised psychodiagnostic evaluation

### 2.2.5 Guidelines for the diagnostic evaluation of ED

	LE	GR
Clinical use of validated questionnaire related to ED may help to assess all sexual function domains and the effect of a specific treatment modality.	3	B
Physical examination is needed in the initial assessment of men with ED to identify underlying medical conditions that may be associated with ED.	4	B
Routine laboratory tests, including glucose-lipid profile and total testosterone, are required to identify and treat any reversible risk factors and lifestyle factors that can be modified.	4	B
Specific diagnostic tests are indicated by only a few conditions.	4	B

### 2.2.6 References

- Davis-Joseph B, Tiefer L, Melman A. Accuracy of the initial history and physical examination to establish the etiology of erectile dysfunction. *Urology* 1995 Mar;45(3):498-502.  
<http://www.ncbi.nlm.nih.gov/pubmed/7879338>
- Hatzichristou D, Hatzimouratidis K, Bekas M, et al. Diagnostic steps in the evaluation of patients with erectile dysfunction. *J Urol* 2002 Aug;168(2):615-20.  
<http://www.ncbi.nlm.nih.gov/pubmed/12131320>
- Lewis RW. Epidemiology of erectile dysfunction. *Urol Clin North Am* 2001 May;28(2):209-16, vii.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402575>
- Rosen RC, Riley A, Wagner G, et al. The international index of erectile function (IIEF): a multidimensional scale for assessment of erectile dysfunction. *Urology* 1997 Jun;49(6):822-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/9187685>
- Mulhall JP, Goldstein I, Bushmakin AG, et al. Validation of the erection hardness score. *J Sex Med* 2007 Nov;4(6):1626-34.  
<http://www.ncbi.nlm.nih.gov/pubmed/17888069>
- Whooley MA, Avins AL, Miranda J, et al. Case-finding instruments for depression. Two questions are as good as many. *J Gen Intern Med* 1997 Jul;12(7):439-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/9229283>

7. Zitzmann M, Faber S, Nieschlag E. Association of specific symptoms and metabolic risks with serum testosterone in older men. *J Clin Endocrinol Metab* 2006 Nov;91(11):4335-43.  
<http://www.ncbi.nlm.nih.gov/pubmed/16926258>
8. O'Connor Db, Lee DM, Corona G, et al. The relationship between sex hormones and sexual function in middle-age and older European men. *J Clin Endocrinol Metab* 2011 Oct;96(10):E1577-87.  
<http://www.ncbi.nlm.nih.gov/pubmed/21849522>
9. Heidenreich A, Bellmunt J, Bolla M, et al. European Association of Urology. EAU guidelines on prostate cancer. Part 1: screening, diagnosis, and treatment of clinically localised disease. *Eur Urol* 2011 Jan;59(1):61-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/21056534>
10. Morales A, Heaton JP. Hormonal erectile dysfunction. Evaluation and management. *Urol Clin North Am* 2001 May;28(2):279-88.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402581>
11. Lue TF, Giuliano F, Montorsi F, et al. Summary of the recommendations on sexual dysfunctions in men. *J Sex Med* 2004 Jul;1(1):6-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422979>
12. Ghanem HM, Salonia A, Martin-Morales A. SOP: Physical Examination and Laboratory Testing for Men with Erectile Dysfunction. *J Sex Med*. 2012 Apr 23. [Epub ahead of print]  
<http://www.ncbi.nlm.nih.gov/pubmed/22524416>
13. Laumann EO, Paik A, Rosen RC. The epidemiology of erectile dysfunction: results from the National Health and Social Life Survey. *Int J Impot Res* 1999 Sep;11(Suppl 1):S60-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/10554933>
14. Gazzaruso C, Coppola A, Montalcini T, et al. Erectile dysfunction can improve the effectiveness of the current guidelines for the screening for asymptomatic coronary artery disease in diabetes. *Endocrine* 2011 Oct;40(2):273-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/21861245>
15. Batty GD, Li Q, Czernichow S, et al. Erectile dysfunction and later cardiovascular disease in men with type 2 diabetes: prospective cohort study based on the ADVANCE (Action in Diabetes and Vascular Disease: Preterax and Diamicon Modified-Release Controlled Evaluation) trial. *J Am Coll Cardiol* 2010 Nov 30;56(23):1908-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/21109113>
16. Dong JY, Zhang YH, Qin LQ. Erectile dysfunction and risk of cardiovascular disease: meta-analysis of prospective cohort studies. *J Am Coll Cardiol* 2011 Sep 20;58(13):1378-85.  
<http://www.ncbi.nlm.nih.gov/pubmed/21920268>
17. DeBusk R, Drory Y, Goldstein I, et al. Management of sexual dysfunction in patients with cardiovascular disease: recommendations of the Princeton Consensus Panel. *Am J Cardiol* 2000 Jul;86(2):175-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/10913479>
18. Kostis J, Jackson G, Rosen R, et al. Sexual dysfunction and cardiac risk (the Second Princeton Consensus Conference). *Am J Cardiol* 2005 Jul;96(2):313-21.  
<http://www.ncbi.nlm.nih.gov/pubmed/16018863>
19. Nehra A, Jackson G, Miner M, et al. The Princeton III Consensus recommendations for the management of erectile dysfunction and cardiovascular disease. *Mayo Clin Proc* 2012 Aug;87(8):766-78.  
<http://www.ncbi.nlm.nih.gov/pubmed/22862865>
20. Hatzichristou DG, Hatzimouratidis K, Ioannides E, et al. Nocturnal penile tumescence and rigidity monitoring in young potent volunteers: reproducibility, evaluation criteria and the effect of sexual intercourse. *J Urol* 1998 Jun;159(6):1921-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/9598488>
21. Meuleman EJ, Diemont WL. Investigation of erectile dysfunction. Diagnostic testing for vascular factors in erectile dysfunction. *Urol Clin North Am* 1995 Nov;22(4):803-19.  
<http://www.ncbi.nlm.nih.gov/pubmed/7483130>
22. Hatzichristou DG, Hatzimouratidis K, Apostolidis A, et al. Hemodynamic characterization of a functional erection. Arterial and corporeal veno-occlusive function in patients with a positive intracavernosal injection test. *Eur Urol* 1999;36(1):60-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10364657>
23. Wespes E, Schulman C. Venous impotence: pathophysiology, diagnosis and treatment. *J Urol* 1993 May;149(5 Pt 2):1238-45.  
<http://www.ncbi.nlm.nih.gov/pubmed/8479008>



24. Rosen RC, Leiblum SR, Spector IP. Psychologically based treatment for male erectile disorder: a cognitive-interpersonal model. *J Sex Marital Ther* 1994 Summer;20(2):67-85.  
<http://www.ncbi.nlm.nih.gov/pubmed/8035472>
25. Hatzichristou D, Rosen RC, Broderick G, et al. Clinical evaluation and management strategy for sexual dysfunction in men and women. *J Sex Med* 2004 Jul;1(1):49-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422983>

## **3. TREATMENT OF ERECTILE DYSFUNCTION**

### **3.1 Treatment options**


The primary goal in the management strategy of a patient with ED is to determine its etiology and treat it when possible, and not to treat the symptom alone. ED may be associated with modifiable or reversible risk factors, including lifestyle or drug-related factors. These factors may be modified either before, or at the same time as, specific therapies are used.

As a rule, ED can be treated successfully with current treatment options, but cannot be cured. The only exceptions are psychogenic ED, post-traumatic arteriogenic ED in young patients, and hormonal causes (e.g., hypogonadism and hyperprolactinaemia), which potentially can be cured with specific treatment.

Most men with ED will be treated with therapeutic options that are not cause specific. This results in a structured treatment strategy that depends on efficacy, safety, invasiveness and cost, as well as patient preference (1). To properly counsel patients with ED, physicians must be fully informed of all available treatment options. In this context, physician-patient (partner) dialogue is essential throughout the management of ED.

The assessment of treatment options must consider patient and partner satisfaction and other QoL factors as well as efficacy and safety. A treatment algorithm for ED is given in Figure 3.

**Figure 3: Treatment algorithm for ED**



**3.2 Lifestyle management in ED with concomitant risk factors**

The basic work-up of the patient must identify reversible risk factors for ED. Lifestyle changes and risk factor modification must precede or accompany any pharmacological treatment.

The potential benefits of lifestyle changes may be particularly important in individuals with ED and specific comorbid cardiovascular or metabolic disorders, such as diabetes or hypertension (2-4). Besides

improving erectile function, aggressive lifestyle changes may also benefit overall cardiovascular and metabolic health, with recent studies supporting the potential of lifestyle intervention to benefit both ED and overall health (5,6).

Although further studies are needed to clarify the role of lifestyle changes in management of ED and related cardiovascular disease, lifestyle changes can be recommended alone or combined with PDE5 therapy. Some studies have suggested that the therapeutic effects of PDE5Is may be enhanced when other comorbidities or risk factors are aggressively managed (7). A significant improvement can be expected as soon as after 3 months of initiating lifestyle changes (8).

However, these results have yet to be confirmed in well-controlled, long-term studies. As a result of the success of pharmacological therapy for ED, clinicians need to provide specific evidence for the benefits of lifestyle change, and hopefully, future research will show this.

### **3.3 Erectile dysfunction after radical prostatectomy**

Use of proerectile drugs following RP is important in achieving postoperative erectile function. Several trials have shown higher rates of erectile function recovery after RP in patients receiving any drug (therapeutic or prophylactic) for ED. Early compared with delayed erectile rehabilitation brings forward the natural healing time of potency (9).

Historically, the treatment options for postoperative ED have included intracavernous injections (10), urethral microsuppository (11), vacuum device therapy (12), and penile implants (13). Intracavernous injections and penile implants are still suggested as second- and third-line treatments, respectively, when oral compounds are not adequately effective or contraindicated for postoperative patients (Sections 3.6 and 3.7).

The management of post-RP ED has been revolutionised by the advent of PDE5Is, with their demonstrated efficacy, ease of use, good tolerability, excellent safety, and positive impact on QoL. Overall, it must be emphasized that post-RP ED patients are poor responders to PDE5Is. However, PDE5Is are the first-line choice of oral pharmacotherapy for post-RP ED in patients who have undergone nerve-sparing (NS) surgery. The choice of PDE5Is as first-line treatment is controversial because the experience (surgical volume) of the surgeon is a key factor in preserving postoperative erectile function, in addition to patient age and NS technique (14-16). In fact, PDE5Is are most effective in patients who have undergone a rigorous NS procedure, which is more commonly performed by large-volume surgeons (14,15).

Early use of high-dose sildenafil after RP has been suggested to be associated with preservation of smooth muscle within the corpora cavernosa (17). Daily sildenafil also results in a greater return of spontaneous normal erectile function after RP compared to placebo following bilateral NSRP in patients who were fully potent before surgery (18,19). The response rate to sildenafil treatment for ED after RP in different trials has ranged from 35% to 75% among those who underwent NSRP and from 0% to 15% among those who underwent non-NSRP (18-21).

Effectiveness of tadalafil and vardenafil as on-demand treatment has been evaluated in post-RP ED.

- A large multicentre trial in Europe and USA has studied tadalafil in patients with ED following bilateral NS surgery. Erectile function was improved in 71% of patients treated with 20 mg tadalafil versus 24% of those treated with placebo, while the rate of successful intercourse attempts was 52% with 20 mg tadalafil versus 26% with placebo (22).
- Similarly, vardenafil has been tested in patients with ED following either unilateral or bilateral NS surgery in a randomised, multicentre, prospective, placebo-controlled study in North America (23). Following bilateral NSRP, erectile function improved by 71% and 60% with 10 and 20 mg vardenafil, respectively. An extended analysis of the same patients undergoing NSRP has underlined the benefit of vardenafil compared to placebo regarding intercourse satisfaction, hardness of erection, orgasmic function, and overall satisfaction with sexual experience (24).

A randomised, double-blind, double-dummy, multicentre, parallel-group study in 87 centres across Europe, Canada, South Africa and the USA, compared on-demand and nightly dosing of vardenafil in men with ED following bilateral NSRP. In patients whose IIEF erectile function domain (IIEF-EF) score was  $\geq 26$  before surgery, vardenafil was efficacious when used on demand, supporting a paradigm shift towards on-demand dosing with PDE5Is in post-RP ED (25). A prospective, randomised, open label, multicentre American study in men with normal erection who underwent bilateral NSRP showed that oral and intraurethral treatment has the same benefit for penile recovery within the first year after surgery (26).

Patients who do not respond to oral PDE5Is after NSRP may be treated with prophylactic intracorporeal alprostadil (27,28). Penile prosthesis remains a satisfactory approach for patients who do not respond to either oral or intracavernous pharmacotherapy or to a vacuum device (29).

### **3.4 Causes of ED that can be potentially treated with a curative intent**

#### **3.4.1 Hormonal causes**

The advice of an endocrinologist may be beneficial for managing patients with hormonal abnormalities. Testosterone deficiency is either a result of primary testicular failure or secondary to pituitary/hypothalamic causes, including a functional pituitary tumour resulting in hyperprolactinaemia.

Testosterone replacement therapy (intramuscular, oral, or transdermal) is effective, but should only be used after other endocrinological causes for testicular failure have been excluded (30). Testosterone replacement is controversial in men with a history of prostate carcinoma (LE: 4) (31). There is limited evidence suggesting that such treatment may not pose an undue risk of prostate cancer recurrence or progression (32). Before initiating testosterone replacement, digital rectal examination, serum PSA test, hematocrit, liver function tests and lipid profile should be performed (33). Patients given androgen therapy should be monitored for clinical response, elevated hematocrit and development of hepatic or prostatic disease. Testosterone therapy is contraindicated in patients with untreated prostate cancer or unstable cardiac disease.

#### **3.4.2 Post-traumatic arteriogenic ED in young patients**

In young patients with pelvic or perineal trauma, surgical penile revascularisation has a 60-70% long-term success rate (34). The lesion must be demonstrated by duplex Doppler study of the penis and confirmed by penile phalloangiography. Corporeal veno-occlusive dysfunction is a contraindication to revascularisation and must be excluded by DICCC. Vascular surgery for veno-occlusive dysfunction is no longer recommended because of poor long-term results (35).

#### **3.4.3 Psychosexual counselling and therapy**

For patients with a significant psychological problem, psychosexual therapy may be given either alone or with another therapeutic approach. Psychosexual therapy requires ongoing follow-up and has had variable results (36).

### **3.5 First-line therapy**

#### **3.5.1 Oral pharmacotherapy**

PDE5 hydrolyses cGMP in the cavernosum tissue. Inhibition of PDE5 results in smooth muscle relaxation with increased arterial blood flow, leading to compression of the subtunical venous plexus and penile erection (37).

Three potent selective PDE5Is have been approved by the European Medicines Agency (EMA) for the treatment of ED. They are not initiators of erection and require sexual stimulation to facilitate an erection.

##### **3.5.1.1 Sildenafil**

Sildenafil was launched in 1998 and was the first PDE5I available on the market. Efficacy is defined as an erection with rigidity sufficient for vaginal penetration. Sildenafil is effective from 30-60 min after administration. Its efficacy is reduced after a heavy, fatty meal due to prolonged absorption. It is administered in doses of 25, 50 and 100 mg. The recommended starting dose is 50 mg and should be adapted according to the patient's response and side effects. Efficacy may be maintained for up to 12 h (38). The pharmacokinetic data of sildenafil are presented in Table 7. Adverse events (Table 8) are generally mild in nature and self-limited by continuous use. The drop-out rate due to adverse events is similar to that with placebo (39).

After 24 weeks in a dose-response study, improved erections were reported by 56%, 77% and 84% of a general ED population taking 25, 50 and 100 mg sildenafil, respectively, compared to 25% of men taking placebo (40). Sildenafil significantly improves patient scores in IIEF, sexual encounter profile (SEP)2, SEP3, and general assessment question (GAQ) and treatment satisfaction.

The efficacy of sildenafil in almost every subgroup of patients with ED has been successfully established. In patients with diabetes, 66.6% reported improved erections (GAQ) and 63% successful intercourse attempts compared to 28.6% and 33% of men taking placebo, respectively (41).

##### **3.5.1.2 Tadalafil**

Tadalafil was licenced for treatment of ED in February 2003 and is effective from 30 min after administration, with peak efficacy after about 2 h. Efficacy is maintained for up to 36 h (42) and is not affected by food. Ten and 20 mg doses have been approved for on-demand treatment of ED. The recommended starting dose is 10 mg and should be adapted according to the patient's response and side effects. Pharmacokinetic data of tadalafil are presented in Table 7. Adverse events (Table 8) are generally mild in nature and self-limited by continuous use. The drop-out rate due to adverse events is similar to that with placebo (43).

In premarketing studies, after 12 weeks of treatment and in a dose-response study, improved erections were reported by 67% and 81% of a general ED population taking 10 and 20 mg tadalafil, respectively, compared to 35% of men in the control placebo group (43). Tadalafil significantly improves patient scores in IIEF, SEP2, SEP3, and GAQ and treatment satisfaction. These results have been confirmed in

postmarketing studies (44).

Tadalafil also improves erections in difficult-to-treat subgroups. In patients with diabetes, 64% reported improved erections (i.e., improved GAQ) versus 25% of patients in the control group, and the change in the final score for IIEF-EF was 7.3 compared to 0.1 for placebo (45). Nevertheless diabetic patients remain poor responders to tadalafil on demand, with a successful intercourse rates increasing from 21.8% with placebo to 45.4 and 49.9% with 10 and 20 mg of tadalafil on demand respectively (46).

### 3.5.1.3 Vardenafil

Vardenafil became commercially available in March 2003 and is effective from 30 min after administration. Its effect is reduced by a heavy, fatty meal (> 57% fat). Five, 10 and 20 mg doses have been approved for on-demand treatment of ED. The recommended starting dose is 10 mg and should be adapted according to the patient's response and side effects. In vitro, it is 10-fold more potent than sildenafil, although this does not necessarily mean greater clinical efficacy (47). Pharmacokinetic data of vardenafil are presented in Table 7. Adverse events (Table 8) are generally mild in nature and self-limited by continuous use, with a drop-out rate similar to that with placebo (48).

After 12 weeks in a dose-response study, improved erections were reported by 66%, 76% and 80% of a general ED population taking 5, 10 and 20 mg vardenafil, respectively, compared with 30% of men taking placebo (49). Vardenafil significantly improved patient scores for IIEF, SEP2, SEP3, and GAQ and treatment satisfaction. Efficacy has been confirmed in postmarketing studies (50).

Vardenafil improves erections in difficult-to-treat subgroups. In patients with diabetes, the final IIEF-EF score was 19 compared to 12.6 for placebo (51). Nevertheless, again, diabetic patients remain poor responders to vardenafil on-demand with a successful intercourse rates increasing from 23% with placebo to 49% and 54 % with 10 and 20 mg of vardenafil on-demand, respectively (51).

Recently, a new formulation of vardenafil has been released, in the form of an orodispersible tablet (ODT). ODT formulations offer improved convenience over film-coated formulations and may be preferred by patients. Absorption is unrelated to food intake and they exhibit better bioavailability compared to film-coated tablets (52). The efficacy of vardenafil ODT has been demonstrated in several randomised controlled trials and did not seem to differ from the regular formulation (53-56).

### 3.5.1.4 Choice or preference between the different PDE5 inhibitors

To date, no data are available from double- or triple-blind multicentre studies comparing the efficacy and/or patient preference for sildenafil, tadalafil, and vardenafil. Choice of drug will depend on the frequency of intercourse (occasional use or regular therapy, 3-4 times weekly) and the patient's personal experience. Patients need to know whether a drug is short- or long-acting, its possible disadvantages, and how to use it.

### 3.5.1.5 On-demand or chronic use of PDE5 inhibitors

Animal studies have shown that chronic use of PDE5Is improves or prevents significantly the intracavernous structure alterations due to age, diabetes, or surgical damage (57-62). No data exists for a human population.

In humans, a randomised study (n = 145) has shown that daily tadalafil led to a significantly higher IIEF-EF score and higher completion of successful intercourse attempts compared to on-demand tadalafil (63). Two major randomised double-blind studies, using 5 and 10 mg/day tadalafil for 12 weeks (n = 268) (64) and 2.5 and 5 mg/day tadalafil for 24 weeks (n = 286) (65), have shown that daily dosing was well tolerated and significantly improved erectile function. However, these studies lacked a comparative on-demand treatment arm. An open-label extension was carried out for both studies in 234 patients for 1 year and 238 patients for 2 years. Tadalafil, 5 mg once daily, was shown to be well tolerated and effective (66). Tadalafil, 5 mg once daily, therefore provides an alternative to on-demand dosing of tadalafil for couples who prefer spontaneous rather than scheduled sexual activities or who anticipate frequent sexual activity, with the advantage that dosing and sexual activity no longer need to be temporally linked. Nevertheless, in the 1-year open-label 5 mg tadalafil extension study followed by 4 weeks wash-out, erectile function was not maintained after discontinuation of therapy in most patients (about 75%).

In 2007, tadalafil 2.5 and 5 mg have been approved by the European Medicines Agency (EMA) for daily treatment of ED. According to EMA, patients who anticipate a frequent use of tadalafil (i.e., at least twice weekly) a once daily regimen with tadalafil 2.5 mg or 5 mg might be considered suitable, based on patient choice and the physician's judgement. In these patients, the recommended dose is 5 mg taken once a day at approximately the same time of day. The dose may be decreased to 2.5 mg once a day based on individual tolerability. The appropriateness of the continuous use of a daily regimen should be reassessed periodically

A double-blind, placebo-controlled, multicentre, parallel-group study was conducted in 236 men with mild-to-moderate ED randomised to receive 10 mg vardenafil once daily plus on-demand placebo for 12 or 24 weeks, or once-daily placebo plus on-demand 10 mg vardenafil for 24 weeks, followed by 4 weeks wash-out

(67). Despite preclinical evidence, the results suggested that once-daily dosing of 10 mg vardenafil does not offer any sustainable effect after cessation of treatment compared to on-demand administration in patients with mild-to-moderate ED.

Other studies (open-label, randomised, crossover studies with limited patient numbers) have shown that chronic, but not on-demand, tadalafil treatment improves endothelial function with a sustained effect after its discontinuation (68, 69). This has been confirmed in another study of chronic sildenafil in men with type 2 diabetes (70).

Recently, in a double-blind, placebo-controlled study of 298 men with diabetes and ED, 2.5 and 5 mg tadalafil once daily for 12 weeks was efficacious and well tolerated. This regimen provides an alternative to on-demand treatment for some men with diabetes (71).

**Table 7: Summary of the key pharmacokinetic data for the three PDE5 inhibitors used to treat ED\***

Parameter	Sildenafil, 100 mg	Tadalafil, 20 mg	Vardenafil, 20 mg
$C_{max}$	560 µg/L	378 µg/L	18.7 µg/L
$T_{max}$	0.8-1 h	2 h	0.9 h
$T_{1/2}$	2.6-3.7 h	17.5 h	3.9 h
AUC	1685 µg.h/L	8066 µg.h/L	56.8 µg.h/L
Protein binding	96%	94%	94%
Bioavailability	41%	NA	15%

$C_{max}$ : maximal concentration,  $T_{max}$ : time-to-maximum plasma concentration;  $T_{1/2}$ : plasma elimination halftime; AUC: area under curve or serum concentration time curve.

\* Fasted state, higher recommended dose. Data adapted from EMA statements on product characteristics.

**Table 8: Common adverse events of the three PDE5 inhibitors used to treat ED\***

Adverse event	Sildenafil	Tadalafil	Vardenafil
Headache	12.8%	14.5%	16%
Flushing	10.4%	4.1%	12%
Dyspepsia	4.6%	12.3%	4%
Nasal congestion	1.1%	4.3%	10%
Dizziness	1.2%	2.3%	2%
Abnormal vision	1.9%		< 2%
Back pain		6.5%	
Myalgia		5.7%	

\* Adapted from EMA statements on product characteristics.

Sildenafil: <http://www.emea.europa.eu/humandocs/Humans/EPAR/viagra/viagra.htm>

Tadalafil: <http://www.emea.europa.eu/humandocs/Humans/EPAR/cialis/cialis.htm>

Vardenafil: <http://www.emea.europa.eu/humandocs/Humans/EPAR/levitra/levitra.htm>

### 3.5.1.6 Safety issues for PDE5 inhibitors

#### 3.5.1.6.1 Cardiovascular safety

Clinical trial results and post-marketing data of sildenafil, tadalafil, and vardenafil have demonstrated no increase in myocardial infarction rates in patients receiving PDE5Is, as part of either double-blind, placebo-controlled trials or open-label studies, or compared to expected rates in age-matched male populations.

None of the PDE5Is had an adverse effect on total exercise time or time-to-ischæmia during exercise testing in men with stable angina (72,73). In fact, they may even improve exercise tests. Sildenafil does not alter cardiac contractility, cardiac output or myocardial oxygen consumption according to available evidence. Chronic or on-demand use is well tolerated with a similar safety profile.

#### 3.5.1.6.2 Nitrates are contraindicated with PDE5 inhibitors

Organic nitrates (e.g., nitroglycerine, isosorbide mononitrate, and isosorbide dinitrate) and other nitrate preparations used to treat angina, as well as amyl nitrite or amyl nitrate ("poppers" used for recreation), are absolute contraindications for the use of PDE5Is. They result in cGMP accumulation and unpredictable falls in blood pressure and symptoms of hypotension. The duration of interaction between organic nitrates and PDE5Is depends upon the PDE5I and nitrate used.

If a PDE5I is taken and the patient develops chest pain, nitroglycerine must be withheld for at least 24 h if sildenafil (and probably also vardenafil) is used (half-life, 4 h), and for at least 48 h if tadalafil is used (half-life, 17.5 h).

If a patient develops angina while taking a PDE5I, other agents may be given instead of nitroglycerine until the appropriate time has passed. If nitroglycerine must be reintroduced following administration of a PDEI, the patient should receive it only after an appropriate interval has elapsed, as described above, and under close medical observation.

#### 3.5.1.6.3 Antihypertensive drugs

Co-administration of PDE5Is with antihypertensive agents (angiotensin-converting enzyme inhibitors, angiotensin-receptor blockers, calcium blockers,  $\beta$ -blockers, and diuretics) may result in small additive decreases in blood pressure, which are usually minor. In general, the adverse event profile of a PDE5I is not made worse by a background of antihypertensive medication, even when the patient is taking several antihypertensive agents.

#### 3.5.1.6.4 $\alpha$ -Blocker interactions

All PDE5Is show some interaction with  $\alpha$ -blockers, which under some conditions may result in orthostatic hypotension.

- Sildenafil labelling currently advises that 50 or 100 mg sildenafil should be used with caution in patients taking an  $\alpha$ -blocker (especially doxazosin). Hypotension is more likely to occur within 4 h following treatment with an  $\alpha$ -blocker. A starting dose of 25 mg is recommended.
- Concomitant treatment with vardenafil should only be initiated if the patient has been stabilised on his alpha-blocker therapy.
- Co-administration of vardenafil with tamsulosin is not associated with clinically significant hypotension (74).
- Tadalafil is not recommended in patients taking doxazosin but this is not the case for tamsulosin, 0.4 mg (75).

These interactions are more pronounced when PDE5Is are given to healthy volunteers not previously taking  $\alpha$ -blockers. Therefore, patients should be stable on  $\alpha$ -blocker therapy prior to initiating combined treatment, and that the lowest dose should be started initially of PDE5Is. Further research is needed into the interaction between other PDE5Is and other  $\alpha$ -blockers (e.g., alfuzosin, once-daily), or mixed  $\alpha/\beta$ -blockers (e.g., carvedilol and labetalol).

#### 3.5.1.6.5 Dosage adjustment

Drugs that inhibit the CYP34A pathway will inhibit the metabolic breakdown of PDE5Is. They include ketoconazole, itraconazole, erythromycin, clarithromycin, and HIV protease inhibitors (ritonavir and saquinavir). Such agents may increase blood levels of PDE5Is, so that lower doses of PDE5Is are necessary.

However, other agents, such as rifampin, phenobarbital, phenytoin and carbamazepine, may induce CYP3A4 and enhance the breakdown of PDE5Is, so that higher doses of PDE5Is are required. Severe kidney or hepatic dysfunction may require dose adjustments or warnings.

#### 3.5.1.7 *Management of non-responders to PDE5 inhibitors*

The two main reasons why patients fail to respond to a PDE5I are either incorrect drug use or lack of efficacy of the drug. The management of non-responders depends upon identifying the underlying cause.

##### 3.5.1.7.1 Check that the patient has been using a licensed medication

There is a large black market in PDE5Is. The amount of active drug in these medications varies enormously and it is important to check how and from which source the patient has obtained his medication.

##### 3.6.1.7.2 Check that the medication has been properly prescribed and correctly used

The main reason why patients fail to use their medication correctly is inadequate counselling from their physician. The main ways in which a drug may be incorrectly used are:

- failure to use adequate sexual stimulation;
- failure to use an adequate dose;
- failure to wait an adequate amount of time between taking the medication and attempting sexual intercourse.

Lack of adequate sexual stimulation: PDE5Is depend for their action upon the release of NO by the parasympathetic nerve endings in the erectile tissue of the penis. The usual stimulus for NO release is sexual

stimulation, and without adequate sexual stimulation (and NO release), the drugs cannot work.

Oral PDE5Is take different times to reach maximal plasma concentrations (76,77). Although pharmacological activity is achieved at plasma levels well below the maximal plasma concentration, there will be a period of time following oral ingestion of the medication during which the drug is ineffective. Even though all three drugs have an onset of action in some patients within 30 min of oral ingestion, most patients require a longer delay between taking the medication, with at least 60 min being required for men using sildenafil and vardenafil, and up to 2 h being required for men using tadalafil (78-80).

Absorption of sildenafil can be delayed by a meal, and absorption of vardenafil can be delayed by a fatty meal (81). Absorption of tadalafil is less affected provided there is enough delay between oral ingestion and an attempt at sexual intercourse (77).

It is possible to wait too long after taking medication before attempting sexual intercourse. The half-life of sildenafil and vardenafil is about 4 h, suggesting that the normal window of efficacy is 6-8 h following drug ingestion, although responses following this time period are well recognised. Tadalafil has a longer half-life of ~17.5 h, so the window of efficacy is much longer at ~36 h.

For financial reasons, some physicians may prescribe only lower doses of a drug. It is important to check that the patient has had an adequate trial of the maximal dose of the drug. Data suggest an adequate trial involves at least six attempts with a particular drug (82).

Data from uncontrolled studies suggests patient education can help salvage an apparent non-responder to a PDE5I. After emphasising the importance of dose, timing, and sexual stimulation to the patient, erectile function can be effectively restored following re-administration of the relevant PDE5I (83-85).

One study (84) went further, and in those patients who still did not respond to the PDE5I, a second-line adjustment was instituted. Patients taking tadalafil were advised to wait at least 2 h between oral ingestion and attempting intercourse. Patients taking vardenafil were advised to use the drug only after a fast. In both patient groups, further apparent non-responders were salvaged. No patients using sildenafil were included in this study.

#### 3.5.1.7.3 Possible manoeuvres in patients correctly using a PDE5 inhibitor

When the patient is using an adequate dose of the drug properly and the response is still inadequate, there are several changes that may improve drug efficacy, although the evidence supporting these interventions is limited.

Erectile dysfunction is typically a symptom of an underlying condition, such as diabetes, hypertension, or dyslipidaemia. There is evidence suggesting that, in patients with hypogonadism, normalisation of serum testosterone might improve response to a PDE5I (86). Modification of other risk factors may also be beneficial as discussed in section 3.2.

A randomised trial has suggested that vardenafil benefits non-responders to sildenafil (87), but because of poor study design, the results are considered to overstate the benefits of switching PDE5Is. However, a randomised, open-label, crossover trial comparing sildenafil and tadalafil has indicated that some patients might respond better to one PDE5I than to another (88). According to the IIEF-EF score, 17% of patients had a better response ( $\geq 5$  points) to tadalafil than to sildenafil, while 14% had a better response to sildenafil than tadalafil.

Although these differences might be explained by variation in drug pharmacokinetics, they do raise the possibility that, despite an identical mode of action, switching to a different PDE5I might be helpful.

Two non-randomized trials have suggested that daily dosing with a PDE5I might salvage some non-responders to intermittent dosing. In one trial (89), some men benefited from regular dosing with either vardenafil or tadalafil, while in the other trial (84) daily dosing with tadalafil salvaged some men who had failed to respond to intermittent dosing with a PDE5I.

Currently, there are no randomised trials to support this intervention. Although tadalafil is licensed for daily dosing at 2.5 and 5 mg, neither sildenafil nor vardenafil are licensed for use in this way.

If drug treatment fails, then patients should be offered an alternative therapy such as intracavernosal injection therapy or use of a vacuum erection device.

#### 3.5.2 **Vacuum erection devices**

Vacuum erection devices (VEDs) provide passive engorgement of the corpora cavernosa, together with a constrictor ring placed at the base of the penis to retain blood within the corpora. Thus, erections with these devices are not normal because they do not use physiological erection pathways. Efficacy, in terms of erections satisfactory for intercourse, is as high as 90%, regardless of the cause of ED and satisfaction rates range between 27% and 94% (90). Men with a motivated, interested, and understanding partner report the highest satisfaction rates. Long-term use of VEDs decreases to 50-64% after 2 years (91). Most men who discontinue use of VEDs do so within 3 months.

The commonest adverse events include pain, inability to ejaculate, petechiae, bruising, and



numbness, which occur in < 30% of patients (92). Serious adverse events (skin necrosis) can be avoided if patients remove the constriction ring within 30 min. VEDs are contraindicated in patients with bleeding disorders or on anticoagulant therapy.

VEDs may be the treatment of choice in well-informed older patients with infrequent sexual intercourse and comorbidity requiring non-invasive, drug-free management of ED (90).

### 3.5.3 **Shockwave therapy**

Recently, the use of low-intensity extracorporeal shock wave therapy was proposed as a novel treatment for ED (93). In the first randomised, double-blind, sham-controlled study, it was demonstrated that low-intensity extracorporeal shock wave therapy had a positive short-term clinical and physiological effect on the erectile function of men who respond to oral PDE5Is (94). Moreover, there are preliminary data showing improvement in penile hemodynamics and endothelial function, as well as IIEF-EF domain score in severe ED patients who are poor responders to PDE5Is (95). The feasibility and tolerability of this treatment, coupled with its potential rehabilitative characteristics, make it an attractive new therapeutic option for men with ED. However, current data are limited and clear recommendations cannot be given. Data regarding the mechanism of action of this procedure are still lacking. In a diabetic rat model, low-intensity extracorporeal shock wave therapy ameliorated diabetes mellitus associated ED by promoting regeneration of nNOS-positive nerves, endothelium, and smooth muscle in the penis. These beneficial effects appear to be mediated by recruitment of endogenous mesenchymal stem cells (MSCs) (96).

## 3.6 **Second-line therapy**

Patients not responding to oral drugs may be offered intracavernous injections. Success rate is high (85%) (97,98). Intracavernous administration of vasoactive drugs was the first medical treatment for ED more than 20 years ago (99).

### 3.6.1 **Intracavernous injections**

#### 3.6.1.1 *Alprostadil*

Alprostadil (Caverject<sup>TM</sup>, Edex/Viridal<sup>TM</sup>) was the first and only drug approved for intracavernous treatment of ED (99). Intracavernous alprostadil is most efficacious as monotherapy at a dose of 5-40 µg; although the 40 µg dose is not registered in every European country. The erection appears after 5-15 min and lasts according to the dose injected. An office-training programme (one or two visits) is required for the patient to learn the correct injection process. In cases of limited manual dexterity, the technique may be taught to their partners. The use of an automatic special pen that avoids a view of the needle can resolve fear of penile puncture and simplifies the technique.

Efficacy rates for intracavernous alprostadil of > 70% have been found in general ED populations, as well as in patient subgroups (e.g., diabetes or cardiovascular disease), with reported sexual activity after 94% of the injections and satisfaction rates of 87-93.5% in patients and 86-90.3% in partners (100-102).

Complications of intracavernous alprostadil include penile pain (50% of patients reported pain but pain reported only after 11% of total injections), prolonged erections (5%), priapism (1%), and fibrosis (2%) (103). Pain is usually self-limited after prolonged use. It can be alleviated with the addition of sodium bicarbonate or local anaesthesia (104,105). Cavernal fibrosis (from a small hematoma) usually clears within a few months after temporary discontinuation of the injection program. However, tunical fibrosis suggests early onset of La Peyronie's disease and may indicate stopping intracavernosal injections indefinitely. Systemic side effects are uncommon. The most common is mild hypotension, especially when using higher doses.

Contraindications include men with a history of hypersensitivity to alprostadil, men at risk of priapism, and men with bleeding disorders.

Despite these favourable data, intracavernous pharmacotherapy is associated with high drop-out rates and limited compliance. Drop-out rates of 41-68% have been described (106-108), with most drop outs occurring within the first 2-3 months. In a comparative study, alprostadil monotherapy had the lowest discontinuation rate (27.5%) compared to overall drug combinations (37.6%), with an attrition rate after the first few months of therapy of 10% per year. Reasons for discontinuation included desire for a permanent modality of therapy (29%), lack of a suitable partner (26%), poor response (23%) (especially among early drop-out patients), fear of needles (23%), fear of complications (22%), and lack of spontaneity (21%). Careful counselling of patients during the office-training phase as well as close follow-up is important in addressing patient withdrawal from an intracavernous injection programme (109).

Today, intracavernous pharmacotherapy is considered a second-line treatment. Patients not responding to oral drugs may be offered intracavernous injections with a high success rate of 85%. Most long-term injection users can switch to sildenafil despite underlying pathophysiology (110-112). However, almost one-third of long-term intracavernous injections users who subsequently also responded to sildenafil preferred to continue with an intracavernous injection programme (112,113).

### 3.6.1.2 Combination therapy

Combination therapy enables a patient to take advantage of the different modes of action of the drugs being used, as well as alleviating side effects by using lower doses of each drug.

- Papaverine (20-80 mg) was the first oral drug used for intracavernous injections. It is most commonly used in combination therapy today due to its high incidence of side effects as monotherapy.
- Phentolamine has been used in combination therapy to increase efficacy. As monotherapy, it produces a poor erectile response.
- Sparse data in the literature support the use of other drugs, such as vasoactive intestinal peptide (VIP), NO donors (linsidomine), forskolin, potassium channel openers, moxisylyte or calcitonin gene-related peptide (CGRP), usually combined with the main drugs (114,115). Most combinations are not standardised and some drugs have limited availability worldwide.
- Papaverine (7.5-45 mg) plus phentolamine (0.25-1.5 mg), and papaverine (8-16 mg) plus phentolamine (0.2-0.4 mg) plus alprostadil (10-20 µg), have been widely used with improved efficacy rates, although they have never been licensed for ED (116-118). The triple combination regimen of papaverine, phentolamine and alprostadil has the highest efficacy rates, reaching 92%; this combination has similar side effects as alprostadil monotherapy, but a lower incidence of penile pain due to lower doses of alprostadil. However, fibrosis is more common (5-10%) when papaverine is used (depending on total dose). In addition, mild hepatotoxicity has been reported with papaverine (119).

Despite high efficacy rates, 5-10% of patients do not respond to combination intracavernous injections. The combination of sildenafil with intracavernous injection of the triple combination regimen may salvage as many as 31% of patients who do not respond to the triple combination alone (120). However, combination therapy is associated with an incidence of adverse effects in 33% of patients, including dizziness in 20% of patients. This strategy can be considered in carefully selected patients before proceeding to a penile implant.

### 3.6.1.3 Intraurethral alprostadil

A specific formulation of alprostadil (125-1000 µg) in a medicated pellet (MUSE™) has been approved for use in ED (121). A vascular interaction between the urethra and the corpora cavernosa enables drug transfer between these structures (121). Erections sufficient for intercourse are achieved in 30-65.9% of patients. In clinical practice, only the higher doses (500 and 1000 µg) have been used with low consistency response rates (121-123). The application of a constriction ring at the root of the penis (ACTIS™) may improve efficacy (123,124).

The most common adverse events are local pain (29-41%) and dizziness with possible hypotension (1.9-14%). Penile fibrosis and priapism are very rare (< 1%). Urethral bleeding (5%) and urinary tract infections (0.2%) are adverse events related to the mode of administration.

Efficacy rates are significantly lower than intracavernous pharmacotherapy (125). Intraurethral pharmacotherapy is a second-line therapy and provides an alternative to intracavernous injections in patients who prefer a less-invasive, although less-efficacious treatment.

## 3.7 Third-line therapy (penile prostheses)

The surgical implantation of a penile prosthesis may be considered in patients who do not respond to pharmacotherapy or who prefer a permanent solution to their problem. The two currently available classes of penile implants include inflatable (2- and 3-piece) and malleable devices (126-129).

Most patients prefer the three-piece inflatable devices due to the more “natural” erections obtained. The three-piece inflatable penile include a separate reservoir placed in the abdominal cavity. Three-piece devices provide the best rigidity and the best flaccidity because they will fill every part of the corporal bodies. However, the two-piece inflatable prosthesis can be a viable option among patients who are deemed high risk of complications with reservoir placements. Malleable prostheses result in a firm penis, which may be manually placed in an erect or flaccid state (126-129).

There are two main surgical approaches for penile prosthesis implantation: peno-scrotal and infrapubic (126-129). The penoscrotal approach provides an excellent exposure, it affords proximal crural exposure if necessary, avoids dorsal nerve injury and permits direct visualisation of pump placement. However, with this this approach the reservoir is blindly placed into the retropubic space, which can be a problem in patients with a history of major pelvic surgery (mainly radical cystectomy). The infrapubic approach has the advantage of reservoir placement under direct vision but the implantation of the pump may be more challenging, and patients are at a slightly increased risk of dorsal nerve injury. Revision surgery is associated with decreased outcomes and may be more challenging.

### 3.7.1 Efficacy and satisfaction rates

Prosthesis implantation has one of the highest satisfaction rates (92-100% in patients and 91-95% in partners) among the treatment options for ED based on appropriate consultation (130-137). Mulhall and colleagues

have used the IIEF and the Erectile Dysfunction Index for Treatment Satisfaction (EDITS) at 3-month intervals following implantation of inflatable penile prostheses. There was a continued improvement in scores for the IIEF and EDITS stabilised 9-12 months following surgery. All variables, including erection, ejaculation, orgasm, and overall sexual satisfaction, improved above baseline values at 1 year after surgery. However, at 3 months following surgery, the results were less satisfactory, suggesting that postoperative counselling and encouragement of patients is important to obtain ultimate satisfaction and positive outcomes at 9-12 months (134).

In a long-term multicentre study of the AMS 700CX three-piece inflatable prosthesis, with a median follow-up of 48 months, 79% of patients were using their device at least twice monthly and 88% would recommend the prosthesis to a friend or relative (135). In another multicentre study with 59 months follow-up, at almost 5 years after surgery, 92.5% of patients were using their prosthesis an average of 1.7 times weekly and excellent or satisfactory results were reported by patients and their partners (132).

Increasingly, in patients with favourable prognosis after RP for prostate cancer, the presence of urinary incontinence and sexual dysfunction (primarily ED and orgasmic dysfunction) is leading doctors to the need for global management of both conditions. Based on appropriate clinical and diagnostic assessments of severity of adverse outcomes depending on patient preference, combination surgery for treatment of ED, with the implant of a penile prosthesis, and stress urinary incontinence (male sling or artificial urinary sphincter) is effective and durable and has an established, definitive role to address this problem (138).

### 3.7.2 **Complications**

The two main complications of penile prosthesis implantation are mechanical failure and infection. Several technical modifications of the most commonly used three-piece prosthesis (AMS 700CX/CXR™ and Coloplast Alpha I™) resulted in mechanical failure rates of < 5% after 5 years follow-up (135,139). Careful surgical technique with proper antibiotic prophylaxis against Gram-positive and Gram-negative bacteria reduces infection rates to 2-3% with primary implantation in low-risk patients. The infection rate may be further reduced to 1-2% by implanting an antibiotic-impregnated prosthesis (AMS Inhibizone™) or hydrophilic-coated prosthesis (Coloplast itan™) (140-143).

Higher risk populations include patients undergoing revision surgery, those with impaired host defenses (immunosuppression, diabetes mellitus, spinal cord injury) or those with penile corporal fibrosis (126-129). Although diabetes is considered to be one of the main risk factors for infection, this is not supported by current data (126-129). Infections, as well as erosions, are significantly higher (9%) in patients with spinal cord injuries (9%) (126-129). Infection requires removal of the prosthesis and antibiotic administration. Alternatively, removal of the infected device with immediate replacement with a new prosthesis has been described using a washout protocol with successful salvages achieved in > 80% of cases (144,145). The majority of revisions are secondary to mechanical failure and combined erosion or infection. Overall, 93% of cases are successfully revised, providing functioning penile prosthesis.

### 3.7.3 **Conclusions**

Penile implants are an attractive solution for patients who do not respond to more conservative therapies. There is enough evidence to recommend this approach in patients not responding to less-invasive treatments due to its high efficacy, safety and satisfaction rates.

### 3.8 Guidelines for the treatment of ED

	LE	GR
Lifestyle changes and risk factor modification must precede or accompany ED treatment.	1a	A
Pro-erectile treatments have to be given at the earliest opportunity after RP.	1b	A
When a curable cause of ED is found, it must be treated first.	1b	B
PDE5Is are first-line therapy.	1a	A
Inadequate/incorrect prescription and poor patient education are the main causes of a lack of response to PDE5Is.	3	B
A VED can be used in patients with a stable relationship.	4	C
Intracavernous injection is second-line therapy.	1b	B
Penile implant is third-line therapy.	4	C

### 3.9 References

- Hatzichristou D, Rosen RC, Broderick G, et al. Clinical evaluation and management strategy for sexual dysfunction in men and women. *J Sex Med* 2004 Jun;1(1):49-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422983>
- Derby CA, Mohr BA, Goldstein I, et al. Modifiable risk factors and erectile dysfunction: can lifestyle changes modify risk? *Urology* 2000 Aug;56(2):302-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/10925098>
- Moyad MA, Barada JH, Lue TF, et al. Sexual Medicine Society Nutraceutical Committee. Prevention and treatment of erectile dysfunction using lifestyle changes and dietary supplements: what works and what is worthless, part I. *Urol Clin North Am* 2004 May;31(2):249-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/15123405>
- Moyad MA, Barada JH, Lue TF, et al; Sexual Medicine Society Nutraceutical Committee. Prevention and treatment of erectile dysfunction using lifestyle changes and dietary supplements: what works and what is worthless, part II. *Urol Clin North Am* 2004 May;31(2):259-73.  
<http://www.ncbi.nlm.nih.gov/pubmed/15123406>
- Gupta BP, Murad MH, Clifton MM, et al. The effect of lifestyle modification and cardiovascular risk factor reduction on erectile dysfunction: a systematic review and meta-analysis. *Arch Intern Med* 2011 Nov 14;171(20):1797-803.  
<http://www.ncbi.nlm.nih.gov/pubmed/21911624>
- Esposito K, Giugliano F, Di Palo C, et al. Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled trial. *JAMA* 2004 Jun;291(24):2978-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/15213209>
- Guay AT. Optimizing response to phosphodiesterase therapy: impact of risk-factor management. *J Androl* 2003 Nov-Dec;24(6 Suppl):S59-S62.  
<http://www.ncbi.nlm.nih.gov/pubmed/14581497>
- Maio G, Saraeb S, Marchiori A. Physical activity and PDE5 inhibitors in the treatment of erectile dysfunction: results of a randomized controlled study. *J Sex Med* 2010 Jun;7(6):2201-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/20367777>
- Salonia A, Burnett AL, Graefen M, et al. Prevention and management of postprostatectomy sexual dysfunctions part 2: recovery and preservation of erectile function, sexual desire, and orgasmic function. *Eur Urol* 2012 Aug;62(2):273-86  
<http://www.ncbi.nlm.nih.gov/pubmed/22575910>
- Montorsi F, Guazzoni G, Strambi LF, et al. Recovery of spontaneous erectile function after nervesparing radical retropubic prostatectomy with and without early intracavernous injections of alprostadil: results of a prospective, randomized trial. *J Urol* 1997 Oct;158(4):1408-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/9302132>
- Raina R, Pahlajani G, Agarwal A, et al. The early use of transurethral alprostadil after radical prostatectomy potentially facilitates an earlier return of erectile function and successful sexual activity. *BJU Int* 2007 Dec;100(6):1317-21.  
<http://www.ncbi.nlm.nih.gov/pubmed/17850385>
- Raina R, Agarwal A, Ausmundson S, et al. Early use of vacuum constriction device following radical prostatectomy facilitates early sexual activity and potentially earlier return of erectile function. *Int J Impot Res* 2006 Jan-Feb;18(1):77-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/16107868>

13. Lane BR, Abouassaly R, Angermeier KW, et al. Three-piece inflatable penile prostheses can be safely implanted after radical prostatectomy through a transverse scrotal incision. *Urology* 2007 Sep;70(30):539-42.  
<http://www.ncbi.nlm.nih.gov/pubmed/17686509>
14. Ficarra V, Novara G, Ahlering TE, et al. Systematic review and meta-analysis of studies reporting potency rates after robot-assisted radical prostatectomy. *Eur Urol*. 2012 Sep;62(3):418-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/22749850>
15. Ayyathurai R, Manoharan M, Nieder AM, et al. Factors affecting erectile function after radical retropubic prostatectomy: results from 1620 consecutive patients. *BJU Int* 2008 Apr;101(7):833-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/18190627>
16. Hollenbeck BK, Dunn RL, Wei JT, et al. Determinants of long-term sexual health outcome after radical prostatectomy measured by a validated instrument. *J Urol* 2003 Apr;169(4):1453-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/12629382>
17. Schwartz EJ, Wong P, Graydon RJ. Sildenafil preserves intracorporeal smooth muscle after radical retropubic prostatectomy. *J Urol* 2004 Feb;171(2 Pt 4):771-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/14713808>
18. Padma-Nathan H, McCullough AR, Levine LA, et al, Study Group. Randomized, double-blind, placebo-controlled study of postoperative nightly sildenafil citrate for the prevention of erectile dysfunction after bilateral nerve-sparing radical prostatectomy. *Int J Impot Res* 2008 Sep-Oct;20(5):479-86.  
<http://www.ncbi.nlm.nih.gov/pubmed/18650827>
19. Bannowsky A, Schulze H, van der Horst C, et al. Recovery of erectile function after nerve-sparing radical prostatectomy: improvement with nightly low-dose sildenafil. *BJU Int* 2008 May;101(10):1279-83.  
<http://www.ncbi.nlm.nih.gov/pubmed/18284406>
20. Raina R, Lakin MM, Agarwal A, et al. Efficacy and factors associated with successful outcome of sildenafil citrate use for erectile dysfunction after radical prostatectomy. *Urology* 2004 May;63(5):960-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/15134989>
21. McCullough AR, Levine LA, Padma-Nathan H. Return of nocturnal erections and erectile function after bilateral nerve-sparing radical prostatectomy in men treated nightly with sildenafil citrate: subanalysis of a longitudinal randomized double-blind placebo-controlled trial. *J Sex Med* 2008 Feb;5(2):476-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/18086170>
22. Montorsi F, Nathan HP, McCullough A, et al. Tadalafil in the treatment of erectile dysfunction following bilateral nerve sparing radical retropubic prostatectomy: a randomized, double-blind, placebo controlled trial. *J Urol* 2004 Sep;172(3):1036-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/15311032>
23. Brock G, Nehra A, Lipshultz LI, et al. Safety and efficacy of vardenafil for the treatment of men with erectile dysfunction after radical retropubic prostatectomy. *J Urol* 2003 Oct;170(4 Pt 1):1278-83.  
<http://www.ncbi.nlm.nih.gov/pubmed/14501741>
24. Nehra A, Grantmyre J, Nadel A, et al. Vardenafil improved patient satisfaction with erectile hardness, orgasmic function and sexual experience in men with erectile dysfunction following nerve sparing radical prostatectomy. *J Urol* 2005 Jun;173(6):2067-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/15879836>
25. Montorsi F, Brock G, Lee J, et al. Effect of nightly versus on-demand vardenafil on recovery of erectile function in men following bilateral nerve-sparing radical prostatectomy. *Eur Urol* 2008 Oct;54(4):924-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/18640769>
26. McCullough AR, Hellstrom WG, Wang R, et al. Recovery of erectile function after nerve sparing radical prostatectomy and penile rehabilitation with nightly intraurethral alprostadil versus sildenafil citrate. *J Urol* 2010 Jun;183(6):2451-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/20403617>
27. Mulhall JP, Bella AJ, Briganti A, et al. Erectile function rehabilitation in the radical prostatectomy patient. *J Sex Med*. 2010 Apr;7(4 Pt 2):1687-98.  
<http://www.ncbi.nlm.nih.gov/pubmed/20388165>
28. Pace G, Del Rosso A, Vicentini C. Penile rehabilitation therapy following radical prostatectomy. *Disabil Rehabil*. 2010;32(14):1204-8  
<http://www.ncbi.nlm.nih.gov/pubmed/20156044>

29. Montague DK. Penile prosthesis implantation for end-stage erectile dysfunction after radical prostatectomy. *Rev Urol* 2005;7(Suppl 2):S51-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16985898>
30. Greenstein A, Mobjeesh NJ, Sofer M, et al. Does sildenafil combined with testosterone gel improve erectile dysfunction in hypogonadal men in whom testosterone supplement therapy alone failed? *J Urol* 2005 Feb;173(2):530-2.  
<http://www.ncbi.nlm.nih.gov/pubmed/15643239>
31. Endogenous Hormones and Prostate Cancer Collaborative Group, Roddam AW, Allen NE, Appleby P, Key TJ. Endogenous sex hormones and prostate cancer: a collaborative analysis of 18 prospective studies. *J Natl Cancer Inst* 2008 Feb 6;100(3):170-83.  
<http://www.ncbi.nlm.nih.gov/pubmed/18230794>
32. Morgentaler A. Testosterone therapy in men with prostate cancer: scientific and ethical considerations. *J Urol* 2013, Jan;189(1 Suppl):S26-33.  
<http://www.ncbi.nlm.nih.gov/pubmed/23234627>
33. Morales A, Heaton JP. Hormonal erectile dysfunction. Evaluation and management. *Urol Clin North Am* 2001 May;28(2):279-88.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402581>
34. Rao DS, Donatucci CF. Vasculogenic impotence. Arterial and venous surgery. *Urol Clin North Am* 2001 May;28(2):309-19.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402583>
35. Wespes E, Wildschutz T, Roumeguere T, et al. The place of surgery for vascular impotence in the third millennium. *J Urol* 2003 Oct;170(4 Pt 1):1284-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/14501742>
36. Rosen RC. Psychogenic erectile dysfunction. Classification and management. *Urol Clin North Am* 2001 May;28(2):269-78.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402580>
37. Lue TF. Erectile dysfunction. *N Engl J Med* 2000 Jun;342(24):1802-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/10853004>
38. Moncada I, Jara J, Subirá D, et al. Efficacy of sildenafil citrate at 12 hours after dosing: re-exploring the therapeutic window. *Eur Urol* 2004 Sep;46(3):357-60; discussion 360-1.  
<http://www.ncbi.nlm.nih.gov/pubmed/15306108>
39. Langtry HD, Markham A. Sildenafil: a review of its use in erectile dysfunction. *Drugs* 1999 Jun;57(6):967-89.  
<http://www.ncbi.nlm.nih.gov/pubmed/10400408>
40. Goldstein I, Lue TF, Padma-Nathan H, et al; Sildenafil Study Group. Oral sildenafil in the treatment of erectile dysfunction. 1998. *J Urol* 2002 Feb;167(2 Pt 2):1197-203.  
<http://www.ncbi.nlm.nih.gov/pubmed/11905901>
41. Stuckey BG, Jadzinsky MN, Murphy LJ, et al. Sildenafil citrate for treatment of erectile dysfunction in men with type 1 diabetes: results of a randomized controlled trial. *Diabetes Care* 2003 Feb;26(2):279-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/12547849>
42. Porst H, Padma-Nathan H, Giuliano F, et al. Efficacy of tadalafil for the treatment of erectile dysfunction at 24 and 36 hours after dosing: a randomized controlled trial. *Urology* 2003 Jul;62(1):121-5; discussion 125-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12837435>
43. Brock GB, McMahon CG, Chen KK, et al. Efficacy and safety of tadalafil for the treatment of erectile dysfunction: results of integrated analyses. *J Urol* 2002 Oct;168(4 Pt 1):1332-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12352386>
44. Montorsi F, Verheyden B, Meuleman E, et al. Long-term safety and tolerability of tadalafil in the treatment of erectile dysfunction. *Eur Urol* 2004 Mar;45(3):339-44; discussion 344-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/15036680>
45. Sáenz de Tejada I, Anglin G, Knight JR, et al. Effects of tadalafil on erectile dysfunction in men with diabetes. *Diabetes Care* 2002 Dec;25(12):2159-64.  
<http://www.ncbi.nlm.nih.gov/pubmed/12453954>
46. Fonseca V, Seftel A, Denne J, et al. Impact of diabetes mellitus on the severity of erectile dysfunction and response to treatment: analysis of data from tadalafil clinical trials. *Diabetologia* 2004 Nov;47(11):1914-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/15599697>

47. Bischoff E, Schneider K. A conscious-rabbit model to study vardenafil hydrochloride and other agents that influence penile erection. *Int J Impot Res* 2001 Aug;13(4):230-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/11494080>
48. Keating GM, Scott LJ. Vardenafil: a review of its use in erectile dysfunction. *Drugs* 2003;63(23):2673-703.  
<http://www.ncbi.nlm.nih.gov/pubmed/14636086>
49. Porst H, Rosen R, Padma-Nathan H, et al. The efficacy and tolerability of vardenafil, a new, oral, selective phosphodiesterase type 5 inhibitor, in patients with erectile dysfunction: the first at-home clinical trial. *Int J Impot Res* 2001 Aug;13(4):192-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/11494074>
50. Potempa AJ, Ulbrich E, Bernard I, et al. Efficacy of vardenafil in men with erectile dysfunction: a flexible-dose community practice study. *Eur Urol* 2004 Jul;46(1):73-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/15183550>
51. Goldstein I, Young JM, Fischer J, et al; Vardenafil Diabetes Study Group. Vardenafil, a new phosphodiesterase type 5 inhibitor, in the treatment of erectile dysfunction in men with diabetes: a multicentre double-blind placebo-controlled fixed-dose study. *Diabetes Care* 2003 Mar;26(3):777-83.  
<http://www.ncbi.nlm.nih.gov/pubmed/12610037>
52. Heinig R, Weimann B, Dietrich H, et al Pharmacokinetics of a new orodispersible tablet formulation of vardenafil: results of three clinical trials. *Clin Drug Investig* 2011;31(1):27-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/20925442>
53. Debruyne FM, Gittelman M, Sperling H, et al. Time to onset of action of vardenafil: a retrospective analysis of the pivotal trials for the orodispersible and film-coated tablet formulations. *J Sex Med* 2011 Oct;8(10):2912-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/21883954>
54. Sperling H, Gittelman M, Norenberg C, et al. Efficacy and safety of an orodispersible vardenafil formulation for the treatment of erectile dysfunction in elderly men and those with underlying conditions: an integrated analysis of two pivotal trials. *J Sex Med* 2011 Jan;8(1):261-71.  
<http://www.ncbi.nlm.nih.gov/pubmed/20807322>
55. Gittelman M, McMahon CG, Rodríguez-Rivera JA, et al. The POTENT II randomised trial: efficacy and safety of an orodispersible vardenafil formulation for the treatment of erectile dysfunction. *Int J Clin Pract* 2010 Apr;64(5):594-603.  
<http://www.ncbi.nlm.nih.gov/pubmed/20456213>
56. Sperling H, Debruyne F, Boermans A, et al. The POTENT I randomized trial: efficacy and safety of an orodispersible vardenafil formulation for the treatment of erectile dysfunction. *J Sex Med* 2010 Apr;7(4 Pt 1):1497-507.  
<http://www.ncbi.nlm.nih.gov/pubmed/20233275>
57. Ahn GJ, Yu JY, Choi SM, et al. Chronic administration of phosphodiesterase 5 inhibitor improves erectile and endothelial function in a rat model of diabetes. *Int J Androl* 2005 Oct;28(5):260-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/16128985>
58. Kovanecz I, Rambhatia A, Ferrini MG, et al. Chronic daily tadalafil prevents the corporal fibrosis and veno-occlusive dysfunction that occurs after cavernosal nerve resection. *BJU Int* 2008 Jan;101(2):203-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/17888043>
59. Ferrini MG, Davila HH, Kovanecz I, et al. Vardenafil prevents fibrosis and loss of corporal smooth muscle that occurs after bilateral cavernosal nerve resection in the rat. *Urology* 2006 Aug;68(2):429-35.  
<http://www.ncbi.nlm.nih.gov/pubmed/16904479>
60. Vignozzi L, Filippi S, Morelli A, et al. Effect of chronic tadalafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *J Sex Med* 2006 May;3(3):419-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/16681467>
61. Ferrini MG, Kovanecz I, Sanchez S, et al. Long-term continuous treatment with sildenafil ameliorates aging-related erectile dysfunction and the underlying corporal fibrosis in the rat. *Biol Reprod* 2007 May;76(5):915-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/17287493>
62. Behr-Roussel D, Gorny D, Mevel K, et al. Chronic sildenafil improves erectile function and endothelium-dependent cavernosal relaxation in rats: lack of tachyphylaxis. *Eur Urol* 2005 Jan;47(1):87-91.  
<http://www.ncbi.nlm.nih.gov/pubmed/15582254>

63. McMahon C. Comparison of efficacy, safety and tolerability of on-demand tadalafil and daily dosed tadalafil for the treatment of erectile dysfunction. *J Sex Med* 2005 May;2(3):415-25.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422874>
64. Porst H, Giuliano F, Glina S, et al. Evaluation of the efficacy and safety of once-a-day dosing of tadalafil 5mg and 10mg in the treatment of erectile dysfunction: results of a multicenter, randomized, double-blind, placebo-controlled trial. *Eur Urol* 2006 Aug;50(2):351-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/16766116>
65. Rajfer J, Aliotta PJ, Steidle CP, et al. Tadalafil dosed once a day in men with erectile dysfunction: a randomized, double-blind, placebo-controlled study in the US. *Int J Impot Res* 2006;19(1):95-103.  
<http://www.ncbi.nlm.nih.gov/pubmed/16871272>
66. Porst H, Rajfer J, Casabé A, et al. Long-term safety and efficacy of tadalafil 5 mg dosed once daily in men with erectile dysfunction. *J Sex Med* 2008 Sep;5(9):2160-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/18557812>
67. Zumbé J, Porst H, Sommer F, et al. Comparable efficacy of once-daily versus on-demand vardenafil in men with mild-to-moderate erectile dysfunction: findings of the RESTORE study. *Eur Urol* 2008 Jul;54(1):204-10.  
<http://www.ncbi.nlm.nih.gov/pubmed/18395326>
68. Rosano GM, Aversa A, Vitale C, et al. Chronic treatment with tadalafil improves endothelial function in men with increased cardiovascular risk. *Eur Urol* 2005 Feb;47(2):214-20.  
<http://www.ncbi.nlm.nih.gov/pubmed/15661417>
69. Aversa A, Greco E, Bruzziches R, et al. Relationship between chronic tadalafil administration and improvement of endothelial function in men with erectile dysfunction: a pilot study. *Int J Impot Res* 2007 Mar-Apr;19(2):200-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16943794>
70. Aversa A, Vitale C, Volterrani M, et al. Chronic administration of Sildenafil improves markers of endothelial function in men with Type 2 diabetes. *Diabet Med* 2008 Jan;25(1):37-44.  
<http://www.ncbi.nlm.nih.gov/pubmed/18199130>
71. Hatzichristou D, Gambla M, Rubio-Aurioles E, et al. Efficacy of tadalafil once daily in men with diabetes mellitus and erectile dysfunction. *Diabet Med* 2008 Feb;25(2):138-46.  
<http://www.ncbi.nlm.nih.gov/pubmed/18290855>
72. Kloner RA. Novel phosphodiesterase type 5 inhibitors: assessing hemodynamic effects and safety parameters. *Clin Cardiol* 2004 Apr;27(4 Suppl 1):I20-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/15115192>
73. Thadani U, Smith W, Nash S, et al. The effect of vardenafil, a potent and highly selective phosphodiesterase-5 inhibitor for the treatment of erectile dysfunction, on the cardiovascular response to exercise in patients with coronary artery disease. *J Am Coll Cardiol* 2002 Dec;40(11):2006-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/12475462>
74. Auerbach SM, Gittelman M, Mazzu A, et al. Simultaneous administration of vardenafil and tamsulosin does not induce clinically significant hypotension in patients with benign prostatic hyperplasia. *Urology* 2004 Nov;64(5):998-1003; discussion 1003-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/15533493>
75. Kloner RA, Jackson G, Emmick JT, et al. Interaction between the phosphodiesterase 5 inhibitor, tadalafil and 2 alpha-blockers, doxazosin and tamsulosin in healthy normotensive men. *J Urol* 2004 Nov;172(5 Pt 1):1935-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/15540759>
76. Nichols DJ, Muirhead GJ, Harness JA. Pharmacokinetics of sildenafil citrate after single oral doses in healthy male subjects: absolute bioavailability, food effects and dose proportionality. *Br J Clin Pharmacol* 2002;53(Suppl 1):5S-12S.  
<http://www.ncbi.nlm.nih.gov/pubmed/11879254>
77. Forgue ST, Patterson BE, Bedding AW, et al. Tadalafil pharmacokinetics in healthy subjects. *Br J Clin Pharmacol* 2006 Mar;61(3):280-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/16487221>
78. Padma-Nathan H, Stecher VJ, Sweeney M, et al. Minimal time to successful intercourse after sildenafil citrate: results of a randomized, double-blind, placebo-controlled trial. *Urology* 2003 Sep;62(3):400-3.  
<http://www.ncbi.nlm.nih.gov/pubmed/12946731>
79. Rosen RC, Padma-Nathan H, Shabsigh R, et al. Determining the earliest time within 30 minutes to erectogenic effect after tadalafil 10 and 20 mg: a multicenter, randomized, double-blind, placebocontrolled, at-home study. *J Sex Med* 2004 Sep;1(2):193-200.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422974>



80. Montorsi F, Padma-Nathan H, Buvat J, et al; Vardenafil Study Group. Earliest time to onset of action leading to successful intercourse with vardenafil determined in an at-home setting: a randomized, double-blind, placebo-controlled trial. *J Sex Med* 2004 Sep; 1(2):168-78.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422971>
81. Rajagopalan P, Mazzu A, Xia C, et al. Effect of high-fat breakfast and moderate-fat evening meal on the pharmacokinetics of vardenafil, an oral phosphodiesterase-5 inhibitor for the treatment of erectile dysfunction. *J Clin Pharmacol* 2003 Mar;43(3):260-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/12638394>
82. McCullough AR, Barada JH, Fawzy A, et al. Achieving treatment optimization with sildenafil citrate (Viagra) in patients with erectile dysfunction. *Urology* 2002 Sep;60(2 Suppl 2):28-38.  
<http://www.ncbi.nlm.nih.gov/pubmed/12414331>
83. Hatzichristou D, Moysidis K, Apostolidis A, et al. Sildenafil failures may be due to inadequate patient instructions and follow-up: a study on 100 non-responders. *Eur Urol* 2005 Apr;47(4):518-22.  
<http://www.ncbi.nlm.nih.gov/pubmed/15774252>
84. Hatzimouratidis K, Moysidis K, Bekos A, et al. Treatment strategy for 'non-responders' to tadalafil and vardenafil: a real-life study. *Eur Urol* 2006 Jul;50(1):126-32.  
<http://www.ncbi.nlm.nih.gov/pubmed/16564127>
85. Gruenwald I, Shenfeld O, Chen J, et al. Positive effect of counseling and dose adjustment in patients with erectile dysfunction who failed treatment with sildenafil. *Eur Urol* 2006 Jul;50(1):134-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/16527391>
86. Greco EA, Spera G, Aversa A. Combining testosterone and PDE5 inhibitors in erectile dysfunction: basic rationale and clinical evidences. *Eur Urol* 2006, Nov;50(5):940-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16979814>
87. Carson CC, Hatzichristou DG, Carrier S, et al; Patient Response with Vardenafil in Sildenafil Non-Responders (PROVEN) Study Group. Erectile response with vardenafil in sildenafil nonresponders: a multicentre, double-blind, 12-week, flexible-dose, placebo-controlled erectile dysfunction clinical trial. *BJU Int* 2004 Dec;94(9):1301-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/15610110>
88. Eardley I, Montorsi F, Jackson G, et al. Factors associated with preference for sildenafil citrate and tadalafil for treating erectile dysfunction in men naive to phosphodiesterase 5 inhibitor therapy: post hoc analysis of data from a multicentre, randomized, open-label, crossover study. *BJU Int* 2007 Jul;100(1):122-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/17552960>
89. McMahon C. Efficacy and safety of daily tadalafil in men with erectile dysfunction previously unresponsive to on-demand tadalafil. *J Sex Med* 2004 Nov;1(3):292-300.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422959>
90. Levine LA, Dimitriou RJ. Vacuum constriction and external erection devices in erectile dysfunction. *Urol Clin North Am* 2001 May;28(2):335-41, ix-x.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402585>
91. Cookson MS, Nadig PW. Long-term results with vacuum constriction device. *J Urol* 1993 Feb;149(2):290-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/8426404>
92. Lewis RW, Witherington R. External vacuum therapy for erectile dysfunction: use and results. *World J Urol* 1997;15(1):78-82.  
<http://www.ncbi.nlm.nih.gov/pubmed/9066099>
93. Vardi Y, Appel B, Jacob G, et al. Can low-intensity extracorporeal shockwave therapy improve erectile function? A 6-month follow-up pilot study in patients with organic erectile dysfunction. *Eur Urol*. 2010 Aug;58(2):243-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/20451317>
94. Vardi Y, Appel B, Kilchevsky A, Gruenwald I. Does low intensity extracorporeal shock wave therapy have a physiological effect on erectile function? Short-term results of a randomized, double-blind, sham controlled study. *J Urol*. 2012 May;187(5):1769-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/22425129>
95. Gruenwald I, Appel B, Vardi Y. Low-intensity extracorporeal shock wave therapy--a novel effective treatment for erectile dysfunction in severe ED patients who respond poorly to PDE5 inhibitor therapy. *J Sex Med*. 2012 Jan;9(1):259-64.  
<http://www.ncbi.nlm.nih.gov/pubmed/22008059>

96. Qiu X, Lin G, Xin Z, et al. Effects of Low-Energy Shockwave Therapy on the Erectile Function and Tissue of a Diabetic Rat Model. *J Sex Med.* 2012 Dec 17. doi: 10.1111/jsm.12024. [Epub ahead of print]  
<http://www.ncbi.nlm.nih.gov/pubmed/23253086>
97. Shabsigh R, Padma-Nathan H, Gittleman M, et al. Intracavernous alprostadil alfadex (EDEX/VIRIDAL) is effective and safe in patients with erectile dysfunction after failing sildenafil (Viagra). *Urology* 2000 Apr;55(4):477-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/10736486>
98. Coombs PG, Heck M, Guhring P, Narus J, Mulhall JP. A review of outcomes of an intracavernosal injection therapy programme. *BJU Int* 2012 Dec;110(11):1787-91.  
<http://www.ncbi.nlm.nih.gov/pubmed/22564343>
99. Leungwattanakij S, Flynn V Jr, Hellstrom WJ. Intracavernosal injection and intraurethral therapy for erectile dysfunction. *Urol Clin North Am* 2001 May;28(2):343-54.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402586>
100. Linet OI, Ogrinc FG. Efficacy and safety of intracavernosal alprostadil in men with erectile dysfunction. The Alprostadil Study Group. *N Engl J Med* 1996 Apr;334(14):873-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/8596569>
101. Porst H. The rationale for prostaglandin E1 in erectile failure: a survey of worldwide experience. *J Urol* 1996 Mar;155(3):802-15.  
<http://www.ncbi.nlm.nih.gov/pubmed/8583582>
102. Heaton JP, Lording D, Liu SN, et al. Intracavernosal alprostadil is effective for the treatment of erectile dysfunction in diabetic men. *Int J Impot Res* 2001 Dec;13(6):317-21.  
<http://www.ncbi.nlm.nih.gov/pubmed/11918246>
103. Lakin MM, Montague DK, VanderBrug Medendorp S, et al. Intracavernous injection therapy: analysis of results and complications. *J Urol* 1990 Jun;143(6):1138-41.  
<http://www.ncbi.nlm.nih.gov/pubmed/2342174>
104. Kattan S. Double-blind randomized crossover study comparing intracorporeal prostaglandin E1 with combination of prostaglandin E1 and lidocaine in the treatment of organic impotence. *Urology* 1995 Jun;45(6):1032-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/7771004>
105. Moriel EZ, Rajfer J. Sodium bicarbonate alleviates penile pain induced by intracavernous injections for erectile dysfunction. *J Urol* 1993 May;149(5 Pt 2):1299-300.  
<http://www.ncbi.nlm.nih.gov/pubmed/8386779>
106. Flynn RJ, Williams G. Long-term follow-up of patients with erectile dysfunction commenced on self injection with intracavernosal papaverine with or without phentolamine. *Br J Urol* 1996 Oct;78(4):628-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/8944522>
107. Sundaram CP, Thomas W, Pryor LE, et al. Long-term follow-up of patients receiving injection therapy for erectile dysfunction. *Urology* 1997 Jun;49(6):932-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/9187703>
108. Gupta R, Kirschen J, Barrow RC 2nd, et al. Predictors of success and risk factors for attrition in the use of intracavernous injection. *J Urol* 1997 May;157(5):1681-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/9112505>
109. Vardi Y, Sprecher E, Gruenwald I. Logistic regression and survival analysis of 450 impotent patients treated with injection therapy: long-term dropout parameters. *J Urol* 2000 Feb;163(2):467-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/10647656>
110. Montorsi F, Althof SE, Sweeney M, et al. Treatment satisfaction in patients with erectile dysfunction switching from prostaglandin E(1) intracavernosal injection therapy to oral sildenafil citrate. *Int J Impot Res* 2003 Dec;15(6):444-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/14671665>
111. Raina R, Lakin MM, Agarwal A, et al. Long-term intracavernous therapy responders can potentially switch to sildenafil citrate after radical prostatectomy. *Urology* 2004 Mar;63(3):532-7; discussion 538.  
<http://www.ncbi.nlm.nih.gov/pubmed/15028452>
112. Hatzichristou DG, Apostolidis A, Tzortzis V, et al. Sildenafil versus intracavernous injection therapy: efficacy and preference in patients on intracavernous injection for more than 1 year. *J Urol* 2000 Oct;164(4):1197-200.  
<http://www.ncbi.nlm.nih.gov/pubmed/10992365>

113. Buvat J, Lemaire A, Ratajczyk J. Acceptance, efficacy and preference of sildenafil in patients on long term auto-intracavernosal therapy: a study with follow-up at one year. *Int J Impot Res* 2002 Dec;14(6): 483-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/12494282>
114. Mulhall JP, Daller M, Traish AM, et al. Intracavernosal forskolin: role in management of vasculogenic impotence resistant to standard 3-agent pharmacotherapy. *J Urol* 1997 Nov;158(5):1752-8; discussion 1758-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/9334594>
115. Buvat J, Costa P, Morlier D, et al. Double-blind multicentre study comparing alprostadil alphacyclodextrin with moxisylyte chlorhydrate in patients with chronic erectile dysfunction. *J Urol* 1998 Jan;159(1):116-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/9400450>
116. Bechara A, Casabé A, Chéliz G, et al. Comparative study of papaverine plus phentolamine versus prostaglandin E1 in erectile dysfunction. *J Urol* 1997 Jun;157(6):2132-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/9146599>
117. Bennett AH, Carpenter AJ, Barada JH. An improved vasoactive drug combination for a pharmacological erection program. *J Urol* 1991 Dec;146(6):1564-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/1719248>
118. McMahon CG. A comparison of the response to the intracavernosal injection of papaverine and phentolamine, prostaglandin E1 and a combination of all three agents in the management of impotence. *Int J Impot Res* 1991;3:113-21.
119. Levine SB, Althof SE, Turner LA, et al. Side effects of self administration of intracavernous papaverine and phentolamine for the treatment of impotence. *J Urol* 1989 Jan;141(1):54-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/2908954>
120. McMahon CG, Samali R, Johnson H. Treatment of intracorporeal injection nonresponse with sildenafil alone or in combination with triple agent intracorporeal injection therapy. *J Urol* 1999 Dec;162(6): 1992-7; discussion 1997-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/10569554>
121. Padma-Nathan H, Hellstrom WJ, Kaiser FE, et al. Treatment of men with erectile dysfunction with transurethral alprostadil. Medicated Urethral System for Erection (MUSE) Study Group. *N Engl J Med* 1997 Jan;336(1):1-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/8970933>
122. Fulgham PF, Cochran JS, Denman JL, et al. Disappointing initial results with transurethral alprostadil for erectile dysfunction in a urology practice setting. *J Urol* 1998 Dec;160(6 Pt 1):2041-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/9817319>
123. Mulhall JP, Jahoda AE, Ahmed A, et al. Analysis of the consistency of intraurethral prostaglandin E(1) (MUSE) during at-home use. *Urology* 2001 Aug;58(2):262-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/11489714>
124. Costa P, Potempa AJ. Intraurethral alprostadil for erectile dysfunction: a review of the literature. *Drugs* 2012 Dec 3;72(17):2243-54.  
<http://www.ncbi.nlm.nih.gov/pubmed/23170913>
125. Shabsigh R, Padma-Nathan H, Gittleman M, et al. Intracavernous alprostadil alfadex is more efficacious, better tolerated, and preferred over intraurethral alprostadil plus optional actis: a comparative, randomized, crossover, multicentre study. *Urology* 2000 Jan;55(4):109-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/10654905>
126. Mulcahy JJ, Austoni E, Barada JH, et al. The penile implant for erectile dysfunction. *J Sex Med* 2004, Jul;1(1):98-109.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422990>
127. Montague DK, Angermeier KW. Penile prosthesis implantation. *Urol Clin North Am* 2001 May;28(2): 355-61, x.  
<http://www.ncbi.nlm.nih.gov/pubmed/11402587>
128. Montague DK. Penile prosthesis implantation in the era of medical treatment for erectile dysfunction. *Urol Clin North Am* 2011, May;38(2):217-2.  
<http://www.ncbi.nlm.nih.gov/pubmed/21621088>
129. Martinez-Salamanca JI, Mueller A, Moncada I, et al. Penile prosthesis surgery in patients with corporal fibrosis: a state of the art review. *J Sex Med* 2011, Jul;8(7):1880-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/21492405>
130. Holloway FB, Farah RN. Intermediate term assessment of the reliability, function and patient satisfaction with the AMS700 Ultrex penile prosthesis. *J Urol* 1997 May;157(5):1687-91.  
<http://www.ncbi.nlm.nih.gov/pubmed/9112506>

131. Tefilli MV, Dubocq F, Rajpurkar A, et al. Assessment of psychosexual adjustment after insertion of inflatable penile prosthesis. *Urology* 1998 Dec;52(6):1106-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/9836564>
132. Montorsi F, Rigatti P, Carmignani G, et al. AMS three-piece inflatable implants for erectile dysfunction: a long-term multi- institutional study in 200 consecutive patients. *Eur Urol* 2000 Jan;37(1):50-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/10671785>
133. Lux M, Reyes-Vallejo L, Morgentaler A, et al. Outcomes and satisfaction rates for the redesigned 2-piece penile prosthesis. *J Urol.* 2007 Jan;177(1):262-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/17162061>
134. Mulhall JP, Ahmed A, Branch J, et al. Serial assessment of efficacy and satisfaction profiles following penile prosthesis surgery. *J Urol* 2003, Apr;169(4):1429-33.  
<http://www.ncbi.nlm.nih.gov/pubmed/12629377>
135. Carson CC, Mulcahy JJ, Govier FE. Efficacy, safety and patient satisfaction outcomes of the AMS 700CX inflatable penile prosthesis: results of a long-term multicenter study. AMS 700CX Study Group. *J Urol* 2000, Aug;164(2):376-80.  
<http://www.ncbi.nlm.nih.gov/pubmed/10893589>
136. Natali A, Olianias R, Fisch M. Penile implantation in Europe: successes and complications with 253 implants in Italy and Germany. *J Sex Med.* 2008 Jun;5(6):1503-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/18410306>
137. Bernal RM, Henry GD. Contemporary patient satisfaction rates for three-piece inflatable penile prostheses. *Adv Urol.* 2012;2012:707321.  
<http://www.ncbi.nlm.nih.gov/pubmed/22899909>
138. Lee D, Westney OL, Wang R. Combination surgery for erectile dysfunction and male incontinence. *Curr Urol Rep* 2011, Dec;12(6):461-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/21956147>
139. Wilson SK, Cleves MA, Delk JR 2nd. Comparison of mechanical reliability of original and enhanced Mentor Alpha I penile prosthesis. *J Urol* 1999 Sep;162(3 Pt 1):715-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/10458350>
140. Carson CC 3rd. Efficacy of antibiotic impregnation of inflatable penile prostheses in decreasing infection in original implants. *J Urol* 2004 Apr;171(4):1611-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/15017233>
141. Wolter CE, Hellstrom WJ. The hydrophilic-coated inflatable penile prosthesis: 1-year experience. *J Sex Med* 2004 Sep;1(2):221-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/16429621>
142. Carson CC 3rd, Mulcahy JJ, Harsch MR. Long-term infection outcomes after original antibiotic impregnated inflatable penile prosthesis implants: up to 7.7 years of follow-up. *J Urol* 2011 Feb;185(2):614-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/21168870>
143. Serefoglu EC, Mandava SH, Gokce A, et al. Long-Term Revision Rate due to Infection in Hydrophilic-Coated Inflatable Penile Prostheses: 11-Year Follow-up. *J Sex Med.* 2012 Jul 3.  
<http://www.ncbi.nlm.nih.gov/pubmed/22759917>
144. Mulcahy JJ. Long-term experience with salvage of infected penile implants. *J Urol* 2000 Feb; 163(2):481-2.  
<http://www.ncbi.nlm.nih.gov/pubmed/10647660>
145. Henry GD, Donatucci CF, Connors W, et al. An outcomes analysis of over 200 revision surgeries for penile prosthesis implantation: a multicenter study. *J Sex Med* 2012 Jan;9(1):309-15.  
<http://www.ncbi.nlm.nih.gov/pubmed/22082149>

## 4. PREMATURE EJACULATION

### 4.1 Introduction

Although premature ejaculation is a very common male sexual dysfunction, it is poorly understood. Patients are often unwilling to discuss their symptoms and many physicians do not know about effective treatments. As a result, patients may be misdiagnosed or mistreated (1). In addition, there is currently no registered pharmacological treatment for PE.

These guidelines provide an evidence-based analysis (2) of published data on definition, clinical evaluation and treatment. It provides recommendations to clinicians on the diagnosis and treatment of PE, without pre-empting physician judgement on individual cases.

### 4.2 Definition of PE

#### 4.2.1 Overview

There have previously been two official definitions of PE, neither of which were universally accepted:

- In the Diagnostic and Statistical Manual of Mental Disorders IV-Text Revision (DSM-IV-TR), PE is defined as a '*persistent or recurrent ejaculation with minimal sexual stimulation before, on, or shortly after penetration and before the person wishes it. The clinician must take into account factors that affect duration of the excitement phase, such as age, novelty of the sexual partner or situation, and recent frequency of sexual activity*' (3).
- In the World Health Organization's International Classification of Diseases-10 (ICD-10), PE is defined as '*the inability to delay ejaculation sufficiently to enjoy lovemaking, which is manifested by either an occurrence of ejaculation before or very soon after the beginning of intercourse (if a time limit is required: before or within 15 seconds of the beginning of intercourse) or ejaculation occurs in the absence of sufficient erection to make intercourse possible. The problem is not the result of prolonged absence from sexual activity*' (4).

Recently, two more definitions have been proposed:

- The Second International Consultation on Sexual and Erectile Dysfunction defined PE as '*ejaculation with minimal stimulation and earlier than desired, before or soon after penetration, which causes bother or distress, and over which the sufferer has little or no voluntary control*' (5).
- The International Society for Sexual Medicine (ISSM) has adopted a completely new definition of PE which is the first evidence-based definition, '*Premature ejaculation is a male sexual dysfunction characterised by ejaculation which always or nearly always occurs prior to or within about one minute of vaginal penetration; and inability to delay ejaculation on all or nearly all vaginal penetrations; and negative personal consequences, such as distress, bother, frustration and/or the avoidance of sexual intimacy*'. It must be noted that this definition is limited to men with lifelong PE who engage in vaginal intercourse since there are insufficient objective data to propose an evidence-based definition for acquired PE (6).

All four definitions have taken into account the time to ejaculation, the inability to control or delay ejaculation, and negative consequences (bother/distress) from PE. However, the major point of debate is quantifying the time to ejaculation, which is usually described by intravaginal ejaculatory latency time (IELT). Several proposals for updating the definition of PE in the forthcoming DSM-V and ICD-11 have been presented (7-11).

#### 4.2.2 Classifications

Premature ejaculation is classified as 'lifelong' (primary) or 'acquired' (secondary) (12). Lifelong PE is characterised by onset from the first sexual experience, remains so during life and ejaculation occurs too fast (before vaginal penetration or < 1-2 min after). Acquired PE is characterised by a gradual or sudden onset following normal ejaculation experiences before onset and time to ejaculation is short (usually not as short as in lifelong PE).

Recently, two more PE syndromes have been proposed (11):

- 'Natural variable PE' is characterised by inconsistent and irregular early ejaculations, representing a normal variation in sexual performance.
- 'Premature-like ejaculatory dysfunction' is characterised by subjective perception of consistent or inconsistent rapid ejaculation during intercourse, while ejaculation latency time is in the normal range or can even last longer. It should not be regarded as a symptom or manifestation of true medical pathology.

The addition of these new types may aid patient stratification, diagnosis and treatment, but their exact role remains to be defined (13).

## 4.3 Epidemiology

### 4.3.1 Prevalence

The major problem in assessing the prevalence of PE is the lack of an accurate (validated) definition at the time the surveys were conducted (14). However, epidemiological research has consistently shown that PE, at least according to the DSM-IV definition, is the most common male sexual dysfunction, with prevalence rates of 20-30% (15-17).

The highest prevalence rate of 31% (men aged 18-59 years) was found by the USA NHLS study (16). Prevalence rates from 18 to 29 years, 30 to 39 years, 40 to 49 years and 50 to 59 years were 30%, 32%, 28% and 55%, respectively. These high prevalence rates may be a result of the dichotomous scale (yes/no) in a single question asking if ejaculation occurred too early, as the prevalence rates in European studies have been significantly lower. A British mailed questionnaire survey estimated that the prevalence rate of PE was between 14% (3 months) and 31% (life-time) (18). A French telephone survey of men aged 18 to 69 years estimated the life-time prevalence of early ejaculation at 15%, including 5% who often had experienced ejaculation prior to penetration and 10% who often had ejaculated too rapidly after vaginal intromission (19). A Swedish interview reported an overall prevalence rate of 9% in men aged 18 to 74 years (20), with prevalence by age being 4% for 18-24 years, 7% for 25-34 years, 8% for 35-49 years, 8% for 50-65 years and 14% for 66-74 years. A Danish study about sexual problems using a questionnaire (12 questions) and an interview (23 questions) reported the prevalence rate for PE to be 14% in men aged 51 years (21). An Italian questionnaire survey recorded a prevalence rate of 21% (22). Finally, in a self-administered questionnaire survey in the Netherlands, the prevalence rate was 13% in men aged 50-78 years (23).

The prevalence of PE in the Premature Ejaculation Prevalence and Attitudes (PEPA) survey (a multinational, internet-based survey) was 22.7% (24.0% in the USA, 20.3% in Germany, and 20.0% in Italy) (17). The Global Study of Sexual Attitudes and Behaviors (GSSAB) survey was conducted in men between 40 and 80 years old in 29 different countries using personal and telephone interviews and self-completed mailed questionnaires; it confirmed that the worldwide prevalence of PE was almost 30%. Except for a low reported rate of PE in Middle Eastern countries (10-15%), prevalence was relatively similar throughout the rest of the world (15). Finally, the prevalence rate of PE was 18% in a five-country European Observational study using the IELT and the Premature Ejaculation Profile (PEP) (24), comparable to those obtained in a similarly designed USA observational study (25).

Further research is needed on the prevalence of lifelong and acquired PE. Limited data suggests that the prevalence of lifelong PE, defined as IELT < 1-2 min, is about 2-5% (20, 25). These results are supported by the moderate genetic influence on PE (26) and low prevalence rates of IELT < 1 min (27).

### 4.3.2 Pathophysiology and risk factors

The aetiology of PE is unknown, with little data to support suggested biological and psychological hypotheses, including anxiety, penile hypersensitivity, and 5-HT receptor dysfunction (5). In addition, the pathophysiology of PE is largely unknown. In contrast to ED, there is no impairment of the physiological events leading up to the forceful expulsion of sperm at the urethral meatus.

A significant proportion of men with ED also experience PE (15). High levels of performance anxiety related to ED may worsen PE, with a risk of misdiagnosing PE instead of the underlying ED.

According to the NHLS, the prevalence of PE is not affected by age (16,17), unlike ED, which increases with age. Premature ejaculation is not affected by marital or income status (16). However, PE is more common in blacks, Hispanic men and men from Islamic backgrounds (28,29) and may be higher in men with a lower educational level (15,16). Other risk factors may include a genetic predisposition (30), poor overall health status and obesity (16), prostate inflammation (31,32), thyroid hormone disorders (33), emotional problems and stress (16,34), and traumatic sexual experiences (1516).

In the only published study on risk modification/prevention strategies (35), successful eradication of causative organisms in patients with chronic prostatitis and PE produced marked improvements in IELT and ejaculatory control compared to untreated patients.

## 4.4 Impact on quality of life

Men with PE are more likely to report low satisfaction with their sexual relationship, low satisfaction with sexual intercourse, difficulty relaxing during intercourse, and less frequent intercourse (36,37). However, the negative impact of PE extends beyond sexual dysfunction. PE has a detrimental effect on self-confidence and the relationship with the partner, and may cause mental distress, anxiety, embarrassment and depression (36,38). Sex drive and overall interest in sex does not appear to be affected by PE (39). However, the partner's satisfaction with the sexual relationship decreased with increasing severity of the man's condition (40).

Despite the serious psychological and QoL consequences of PE, few men seek treatment. In the GSSAB survey, 78% of men who self-reported a sexual dysfunction sought no professional help or advice for their sexual problems (15), with men more likely to seek treatment for ED than for PE (15). In the PEPA survey,

only 9% of men with self-reported PE consulted a doctor (17).

The main reasons for not discussing PE with their physician are patient embarrassment and a belief that there is no treatment. Physicians are often uncomfortable discussing sexuality with their patients usually because of embarrassment and a lack of training or expertise in treating PE (41,42). Physicians need to encourage patients to talk about PE.

#### 4.5 Diagnosis

Diagnosis of PE is based on the patient's medical and sexual history (43,44). History should classify PE as lifelong or acquired and determine whether PE is situational (under specific circumstances or with a specific partner) or consistent. Special attention should be given to the duration time of ejaculation, degree of sexual stimulus, impact on sexual activity and QoL, and drug use or abuse. It is also important to distinguish PE from ED.

Many patients with ED develop secondary PE caused by the anxiety associated with difficulty in attaining and maintaining an erection (45). Furthermore, some patients are not aware that loss of erection after ejaculation is normal and may erroneously complain of ED, while the actual problem is PE (46).

There are several overlapping definitions of PE, with four shared factors (Table 7), resulting in a multidimensional diagnosis (47).

**Table 9: Common factors in different definitions of ED**

Time to ejaculation assessed by IELT
Perceived control
Distress
Interpersonal difficulty related to the ejaculatory dysfunction

##### 4.5.1 Intravaginal ejaculatory latency time (IELT)

The use of IELT alone is not sufficient to define PE, as there is significant overlap between men with and without PE (24,25). Moreover, IELT has a significant direct effect on perceived control over ejaculation, but not a significant direct effect on ejaculation-related personal distress or satisfaction with sexual intercourse (48). In addition, perceived control over ejaculation has a significant direct effect on both ejaculation-related personal distress and satisfaction with sexual intercourse (each showing direct effects on interpersonal difficulty related to ejaculation).

In everyday clinical practice, self-estimated IELT is sufficient. Self-estimated and stopwatch-measured IELT are interchangeable and correctly assign PE status with 80% sensitivity and 80% specificity (49). Specificity can be improved further to 96% by combining IELT with a single-item patient-reported outcome (PRO) on control over ejaculation and satisfaction with sexual intercourse (scale ranging from 0 = very poor to 4 = very good) and on personal distress and interpersonal difficulty (0 = not at all to 4 = extremely). However, stopwatch-measured IELT is necessary in clinical trials.

##### 4.5.2 PE assessment questionnaires

The need to assess PE objectively has led to the development of several questionnaires based on the use of PROs (47). Only two questionnaires can discriminate between patients who have PE and those who do not:

- Premature Ejaculation Diagnostic Tool (PEDT): five-item questionnaire based on focus groups and interviews from the USA, Germany and Spain. Assesses control, frequency, minimal stimulation, distress and interpersonal difficulty (50,51).
- Arabic Index of Premature Ejaculation (AIPE): seven-item questionnaire developed in Saudi Arabia assesses sexual desire, hard erections for sufficient intercourse, time to ejaculation, control, satisfaction for the patient and partner, anxiety or depression (52).

These tools are a significant step in simplifying the methodology of PE drug studies, though further cross-cultural validation is needed (53).

Other questionnaires used to characterise PE and determine treatment effects include the PEP (25), Index of Premature Ejaculation (IPE), (54) and Male Sexual Health Questionnaire Ejaculatory Dysfunction (MSHQ-EjD) (55). Currently, their role is optional in everyday clinical practice.

##### 4.5.3 Physical examination and investigations

Physical examination is part of the initial assessment of men with PE. It includes a brief examination of the vascular, endocrine and neurological systems to identify underlying medical conditions associated with PE or other sexual dysfunctions, such as chronic illness, endocrinopathy, autonomic neuropathy, Peyronie's disease,

urethritis or prostatitis. Laboratory or physiological testing should be directed by specific findings from history or physical examination and is not routinely recommended (44).

#### 4.6 Recommendations on the diagnosis of PE

	LE	GR
Diagnosis and classification of PE is based on medical and sexual history. It should be multidimensional and assess IELT, perceived control, distress and interpersonal difficulty due to the ejaculatory dysfunction.	1a	A
Clinical use of self-estimated IELT is adequate. Stopwatch-measured IELT is necessary in clinical trials.	2a	B
Patient-reported outcomes (PROs) have the potential to identify men with PE. Further research is needed before PROs can be recommended for clinical use.	3	C
Physical examination may be necessary in initial assessment of PE to identify underlying medical conditions that may be associated with PE or other sexual dysfunctions, particularly ED.	3	C
Routine laboratory or neurophysiological tests are not recommended. They should only be directed by specific findings from history or physical examination.	3	C

#### 4.7 References

- Rosenberg MT, Sadovsky R. Identification and diagnosis of premature ejaculation. *Int J Clin Pract* 2007 Jun;61(6):903-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/17504352>
- Oxford Centre for Evidence-Based Medicine Levels of Evidence (May 2009). Produced by Bob Phillips, Chris Ball, Dave Sackett, Doug Badenoch, Sharon Straus, Brian Haynes, Martin Dawes since November 1998. Updated by Jeremy Howick March 2009.  
<http://www.cebm.net/index.aspx?o=1025> [Access date January 2013]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th Ed. Text Revision. Washington, D.C., American Psychiatric Publishing, Inc, 2000.
- International Classification of Diseases and Related Health Problems*. 10th Ed. Geneva, World Health Organization, 1994.
- McMahon CG, Abdo C, Incrocci L, et al. Disorders of orgasm and ejaculation in men. *J Sex Med* 2004 Jul;1(1):58-65.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422984>
- McMahon CG, Althof SE, Waldinger MD, et al. An evidence-based definition of lifelong premature ejaculation: report of the International Society for Sexual Medicine (ISSM) ad hoc committee for the definition of premature ejaculation. *J Sex Med* 2008 Jul;5(7):1590-606.  
<http://www.ncbi.nlm.nih.gov/pubmed/18466262>
- Balon R, Segraves RT, Clayton A. Issues for DSM-V: sexual dysfunction, disorder, or variation along normal distribution: toward rethinking DSM criteria of sexual dysfunctions. *Am J Psychiatry* 2007 Feb;164(2):198-200. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/17267778>
- Waldinger MD, Schweitzer DH. The DSM-IV-TR is an inadequate diagnostic tool for premature ejaculation. *J Sex Med* 2007 May;4(3):822-3. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/17498112>
- Waldinger MD, Schweitzer DH. The use of old and recent DSM definitions of premature ejaculation in observational studies: a contribution to the present debate for a new classification of PE in the DSM-V. *J Sex Med* 2008 May;5(5):1079-87.  
<http://www.ncbi.nlm.nih.gov/pubmed/18331260>
- Waldinger MD, Schweitzer DH. Changing paradigms from a historical DSM-III and DSM-IV view toward an evidence-based definition of premature ejaculation. Part I-validity of DSM-IV-TR. *J Sex Med* 2006 Jul;3(4):682-92.  
<http://www.ncbi.nlm.nih.gov/pubmed/16839325>
- Waldinger MD, Schweitzer DH. Changing paradigms from a historical DSM-III and DSM-IV view toward an evidence-based definition of premature ejaculation. Part II-proposals for DSM-V and ICD-11. *J Sex Med* 2006 Jul;3(4):693-705.  
<http://www.ncbi.nlm.nih.gov/pubmed/16839326>



12. Godpodinoff ML. Premature ejaculation: clinical subgroups and etiology. *J Sex Marital Ther* 1989 Summer;15(2):130-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/2769774>
13. Waldinger MD. Premature ejaculation: state of the art. *Urol Clin North Am* 2007 Nov;34(4):591-9, vii-viii.  
<http://www.ncbi.nlm.nih.gov/pubmed/17983899>
14. Waldinger MD. The neurobiological approach to premature ejaculation. *J Urol* 2002 Dec;168(6):2359-67.  
<http://www.ncbi.nlm.nih.gov/pubmed/12441918>
15. Laumann EO, Nicolosi A, Glasser DB, et al; GSSAB Investigators' Group. Sexual problems among women and men aged 40-80 y: prevalence and correlates identified in the Global Study of Sexual Attitudes and Behaviors. *Int J Impot Res* 2005 Jan-Feb;17(1):39-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/15215881>
16. Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999 Feb;281(6):537-44.  
<http://www.ncbi.nlm.nih.gov/pubmed/10022110>
17. Porst H, Montorsi F, Rosen RC, et al. The Premature Ejaculation Prevalence and Attitudes (PEPA) survey: prevalence, comorbidities, and professional help-seeking. *Eur Urol* 2007 Mar;51(5):816-23; discussion 824.  
<http://www.ncbi.nlm.nih.gov/pubmed/16934919>
18. Dunn KM, Croft PR, Hackett GI. Sexual problems: a study of the prevalence and need for health care in the general population. *Fam Pract* 1998 Dec;15(6):519-24.  
<http://www.ncbi.nlm.nih.gov/pubmed/10078790>
19. Spira A, Bajos N, Giami A, et al. Cross-national comparisons of sexual behavior surveys--methodological difficulties and lessons for prevention. *Am J Public Health* 1998 May;88(5):730-1. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/9585733>
20. Fugl-Meyer AR, Sjogren Fugl-Meyer K. Sexual disabilities, problems and satisfaction in 18-74 year old Swedes. *Scan J Sexol* 1999;2:79-105.
21. Solstad K, Hertoft P. Frequency of sexual problems and sexual dysfunction in middle-aged Danish men. *Arch Sex Behav* 1993 Feb;22(1):51-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/8435039>
22. Basile Fasolo C, Mirone V, Gentile V, et al; Andrology Prevention Week centers; Italian Society of Andrology (SIA). Premature ejaculation: prevalence and associated conditions in a sample of 12,558 men attending the andrology prevention week 2001—a study of the Italian Society of Andrology (SIA). *J Sex Med* 2005 May;2(3):376-82.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422869>
23. Blanker MH, Bosch JL, Groeneveld FP, et al. Erectile and ejaculatory dysfunction in a community-based sample of men 50 to 78 years old: prevalence, concern, and relation to sexual activity. *Urology* 2001 Apr;57(4):763-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/11306400>
24. Giuliano F, Patrick DL, Porst H, et al; 3004 Study Group. Premature ejaculation: results from a five-country European observational study. *Eur Urol* 2008 May;53(5):1048-57.  
<http://www.ncbi.nlm.nih.gov/pubmed/17950985>
25. Patrick DL, Althof SE, Pryor JL, et al. Premature ejaculation: an observational study of men and their partners. *J Sex Med* 2005 May;2(3):358-67.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422867>
26. Jern P, Santtila P, Witting K, et al. Premature and delayed ejaculation: genetic and environmental effects in a population-based sample of Finnish twins. *J Sex Med* 2007 Nov;4(6):1739-49.  
<http://www.ncbi.nlm.nih.gov/pubmed/17888070>
27. Waldinger MD, Quinn P, Dilleen M, et al. A multinational population survey of intravaginal ejaculation latency time. *J Sex Med* 2005 Jul;2(4):492-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422843>
28. Richardson D, Goldmeier D. Premature ejaculation—does country of origin tell us anything about etiology? *J Sex Med* 2005 Jul;2(4):508-12.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422845>
29. Carson C, Gunn K. Premature ejaculation: definition and prevalence. *Int J Impot Res* 2006 Sep-Oct;18(Suppl 1):S5-13.  
<http://www.ncbi.nlm.nih.gov/pubmed/16953247>

30. Waldinger MD, Rietschel M, Nöthen MM, et al. Familial occurrence of primary premature ejaculation. *Psychiatr Genet* 1998 Spring;8(1):37-40. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/9564687>
31. Screponi E, Carosa E, Di Stasi SM, et al. Prevalence of chronic prostatitis in men with premature ejaculation. *Urology* 2001 Aug;58(2):198-202.  
<http://www.ncbi.nlm.nih.gov/pubmed/11489699>
32. Shamloul R, el-Nashaar A. Chronic prostatitis in premature ejaculation: a cohort study in 153 men. *J Sex Med* 2006 Jan;3(1):150-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/16409229>
33. Carani C, Isidori AM, Granata A, et al. Multicenter study on the prevalence of sexual symptoms in male hypo- and hyperthyroid patients. *J Clin Endocrinol Metab* 2005 Dec;90(12):6472-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/16204360>
34. Dunn KM, Croft PR, Hackett GI. Association of sexual problems with social, psychological, and physical problems in men and women: a cross sectional population survey. *J Epidemiol Community Health* 1999 Mar;53(3):144-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/10396490>
35. El-Nashaar A, Shamloul R. Antibiotic treatment can delay ejaculation in patients with premature ejaculation and chronic bacterial prostatitis. *J Sex Med* 2007 Mar;4(2):491-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/17367444>
36. Rowland D, Perelman M, Althof S, et al. Self-reported premature ejaculation and aspects of sexual functioning and satisfaction. *J Sex Med* 2004 Sep;1(2):225-32.  
<http://www.ncbi.nlm.nih.gov/pubmed/16429622>
37. Rowland DL, Patrick DL, Rothman M, et al. The psychological burden of premature ejaculation. *J Urol* 2007 Mar;177(3):1065-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/17296413>
38. Symonds T, Roblin D, Hart K, et al. How does premature ejaculation impact a man's life? *J Sex Marital Ther* 2003 Oct-Dec;29(5):361-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/14504007>
39. Riley A, Segraves RT. Treatment of premature ejaculation. *Int J Clin Pract* 2006 Jun;60(6):694-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16805755>
40. Byers ES, Grenier G. Premature or rapid ejaculation: heterosexual couples' perceptions of men's ejaculatory behavior. *Arch Sex Behav* 2003 Jun;32(3):261-70.  
<http://www.ncbi.nlm.nih.gov/pubmed/12807298>
41. Sotomayor M. The burden of premature ejaculation: the patient's perspective. *J Sex Med* 2005 May;2(Suppl 2):110-4.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422797>
42. Solursh DS, Ernst JL, Lewis RW, et al. The human sexuality education of physicians in North American medical schools. *Int J Impot Res* 2003 Oct;15(Suppl 5):S41-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/14551576>
43. Sharlip I. Diagnosis and treatment of premature ejaculation: the physician's perspective. *J Sex Med* 2005 May;2(Suppl 2):103-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422796>
44. Shabsigh R. Diagnosing premature ejaculation: a review. *J Sex Med* 2006 Sep;3(Suppl4):318-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/16939476>
45. Rowland DL, Slob AK. Premature ejaculation: psychophysiological considerations in theory, research, and treatment. *Annu Rev Sex Res* 1997;8:224-53. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/10051895>
46. Althof SE. Prevalence, characteristics and implications of premature ejaculation/rapid ejaculation. *J Urol* 2006 Mar;175(3 Pt 1):842-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/16469562>
47. Althof SE, Symonds T. Patient reported outcomes used in the assessment of premature ejaculation. *Urol Clin North Am* 2007 Nov;34(4):581-9, vii.  
<http://www.ncbi.nlm.nih.gov/pubmed/17983898>
48. Patrick DL, Rowland D, Rothman M. Interrelationships among measures of premature ejaculation: the central role of perceived control. *J Sex Med* 2007 May;4(3):780-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/17419817>
49. Rosen RC, McMahon CG, Niederberger C, et al. Correlates to the clinical diagnosis of premature ejaculation: results from a large observational study of men and their partners. *J Urol* 2007 Mar;177(3):1059-64; discussion 1064.  
<http://www.ncbi.nlm.nih.gov/pubmed/17296411>

50. Symonds T, Perelman M, Althof S, et al. Further evidence of the reliability and validity of the premature ejaculation diagnostic tool. *Int J Impot Res* 2007 Sep-Oct;19(5):521-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/17568761>
51. Symonds T, Perelman MA, Althof S, et al. Development and validation of a premature ejaculation diagnostic tool. *Eur Urol* 2007 Aug;52(2):565-73.  
<http://www.ncbi.nlm.nih.gov/pubmed/17275165>
52. Arafa M, Shamloul R. Development and evaluation of the Arabic Index of Premature Ejaculation (AIPE). *J Sex Med* 2007 Nov;4(6):1750-6.  
<http://www.ncbi.nlm.nih.gov/pubmed/17970977>
53. McMahon CG. Ejaculatory latency vs. patient-reported outcomes (PROs) as study end points in premature ejaculation clinical trials. *Eur Urol* 2007 Aug;52(2):321-3. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/17445975>
54. Althof S, Rosen R, Symonds T, et al. Development and validation of a new questionnaire to assess sexual satisfaction, control, and distress associated with premature ejaculation. *J Sex Med* 2006 May;3(3):465-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/16681472>
55. Rosen RC, Catania JA, Althof SE, et al. Development and validation of four-item version of Male Sexual Health Questionnaire to assess ejaculatory dysfunction. *Urology* 2007 May;69(5):805-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/17482908>

## 4.8 Treatment

In many relationships, PE causes few, if any, problems. In such cases, treatment should be limited to psychosexual counselling. Before beginning treatment, it is essential to discuss patient expectations thoroughly. Erectile dysfunction, in particular, or other sexual dysfunction or genitourinary infection (e.g. prostatitis), should be treated first or at the same time as PE.

Various behavioural techniques have demonstrated benefit in treating PE and are indicated for patients uncomfortable with pharmacological therapy. In lifelong PE, behavioural techniques are not recommended for first-line treatment. They are time-intensive, require the support of a partner and can be difficult to do. In addition, long-term outcomes of behavioural techniques for PE are unknown.

Pharmacotherapy is the basis of treatment in lifelong PE. Since no drug for PE has been approved by the EMEA or FDA, all medical treatments are off-label indications. Only chronic selective serotonin reuptake inhibitors (SSRIs) and on-demand topical anaesthetic agents have consistently shown efficacy in PE. Again, long-term outcomes for pharmacological treatments are unknown.

An evidence-based analysis of all current treatment modalities was performed. Levels of evidence and grade of recommendation are provided and a treatment algorithm is presented (Figure 3).

### 4.8.1 Psychological/behavioural strategies

Behavioural strategies mainly include the 'stop-start' programme developed by Semans (1) and its modification, the 'squeeze' technique, proposed by Masters and Johnson.

- In the 'stop-start' programme, the partner stimulates the penis until the patient feels the urge to ejaculate. At this point, he instructs his partner to stop, waits for the sensation to pass and then stimulation is resumed.
- The 'squeeze' technique is similar but the partner applies manual pressure to the glans just before ejaculation until the patient loses his urge.

Both these procedures are typically applied in a cycle of three pauses before proceeding to orgasm. Behavioural strategies are based on the hypothesis that PE occurs because the man fails to appreciate the sensations of heightened arousal and to recognise the feelings of ejaculatory inevitability. Re-training may attenuate stimulus-response connections by gradually exposing the patient to progressively more intense and more prolonged stimulation, while maintaining the intensity and duration of the stimulus just below the threshold for triggering the response. There are several modifications of these techniques making comparison difficult.

Masturbation before anticipation of sexual intercourse is a technique used by many younger men. Following masturbation, the penis is desensitised resulting in greater ejaculatory delay after the refractory period is over. In a different approach, the man learns to recognise the signs of increased sexual arousal and how to keep his level of sexual excitement below the intensity that elicits the ejaculatory reflex. Efficacy is similar to the 'start-stop' programme (2).

Overall, success rates of 50-60% have been reported short term (3,4). However, there is no controlled research to support the efficacy of behavioural techniques, while a double-blind, randomised, crossover study showed that pharmacological treatment (chlomipramine, sertraline, paroxetine and sildenafil) resulted in greater

IELT prolongation than behavioural therapy (5). Furthermore, clinical experience suggests that improvements achieved with these techniques are generally not maintained long term (6,7).

#### 4.8.1.1 Recommendation on the psychological/behavioural treatment of PE

	LE	GR
Psychological/behavioural therapies may be attempted but no clinical data exists supporting prolonged effect.	3	C

#### 4.8.2 Topical anaesthetic agents

The use of local anaesthetics to delay ejaculation is the oldest form of pharmacological therapy for PE (8). Several trials (9,10) support the hypothesis that topical desensitising agents reduce the sensitivity of the glans penis so delaying ejaculatory latency, but without adversely affecting the sensation of ejaculation.

##### 4.8.2.1 Lidocaine-prilocaine cream

In a randomised, double-blind, placebo-controlled trial, lidocaine-prilocaine cream increased the IELT from 1 min in the placebo group to 6.7 min in the treatment group (11). In another randomised, double-blind, placebo-controlled trial, lidocaine-prilocaine cream significantly increased the stopwatch-measured IELT from 1.49 to 8.45 min while no difference was recorded in the placebo group (1.67 to 1.95 min) (12). Lidocaine-prilocaine cream (5%) is applied for 20 to 30 min prior to intercourse. Prolonged application of topical anaesthetic (30 to 45 min) may result in loss of erection due to numbness of the penis in a significant percentage of men (11). A condom is required to avoid diffusion of the topical anaesthetic agent into the vaginal wall causing numbness in the partner. Alternatively, the condom may be removed prior to sexual intercourse and the penis washed clean of any residual active compound. Although no significant side-effects have been reported, topical anaesthetics are contraindicated in patients or partners with an allergy to any component of the product.

An aerosol formulation of lidocaine 7.5 mg plus prilocaine 2.5 mg (Topical Eutectic Mixture for Premature Ejaculation, TEMPE (13) has been evaluated in a phase II study (14). Intravaginal ejaculatory latency time increased from a baseline of 1 min to 4.9 min in the TEMPE-treated group compared to an increase from baseline of 0.9 min to 1.6 min ( $p < 0.01$ ) in the placebo-treated group. It has been suggested that lidocaine-prilocaine can penetrate the glans within 5–10 min, but penetrates intact keratinised skin less easily, reducing penile numbness and ED (14,15).

Finally, in a randomised, double-blind, placebo-controlled, parallel-group study, lidocaine-prilocaine cream showed similar efficacy to combination with sildenafil (50 mg before coitus) and significantly better efficacy than sildenafil alone (16). However, no specific data on estimated IELT were provided.

##### 4.8.2.2 SS-cream

SS-cream is a topical anaesthetic agent made from the extracts of nine herbs. It is applied to the glans penis 1 h before and washed off immediately prior to coitus. SS-cream increased the vibratory threshold in a dose-dependent fashion, as well as the latency and amplitude of somatosensory-evoked potentials measured at the glans penis (17,18). In a double-blind, randomised, placebo-controlled study (19), application of 0.2 g SS-cream improved IELT from 1.37 min to 10.92 min in the treatment group versus 2.45 min in the placebo group. Sexual satisfaction improved by 82% in the treatment group versus 20% in the placebo group. Mild local burning and mild pain were reported by 18.5% of patients. No adverse effects on sexual function or partner or systemic side-effects were observed.

#### 4.8.2.3 Recommendations on the topical therapy for PE

	LE	GR
Lidocaine-prilocaine cream.	1B	A
SS-cream.	1B	A

#### 4.8.3 Selective serotonin reuptake inhibitors

Ejaculation is mediated by a spinal ejaculation generator (20, 21) and by descending supraspinal modulation from several brain regions. The neurotransmitter 5-hydroxytryptamine (5-HT, serotonin) is also involved in ejaculatory control. The retarding effect of 5-HT on ejaculation is probably due to central activation (i.e. spinally and supraspinally) of 5-HT<sub>1B</sub> and 5-HT<sub>2C</sub> receptors, while stimulation of 5-HT<sub>1A</sub> receptors precipitates ejaculation.

Selective serotonin reuptake inhibitors (SSRIs) are used to treat mood disorders, but can delay

ejaculation and are therefore widely used 'off-label' for PE. As in depression, SSRIs must be given for 1 to 2 weeks to be effective in PE (22). Chronic SSRI administration causes prolonged increases in synaptic cleft serotonin, which desensitise the 5-HT<sub>1A</sub> and 5-HT<sub>1B</sub> receptors (23). Clomipramine, the most serotonergic tricyclic antidepressant, was first reported in 1973 as an effective PE treatment (24). Selective serotonin reuptake inhibitors have revolutionised treatment of PE, but they have also changed our understanding of PE since the first publication on paroxetine in 1970 (25). Today, daily treatment with SSRIs has become the first choice of treatment in PE. Commonly used SSRIs include citalopram, fluoxetine, fluvoxamine, paroxetine and sertraline, all of which have a similar pharmacological mechanism of action.

A systematic review and meta-analysis of all drug treatment studies reported that, despite methodological problems in most studies, there still remained several, well-designed, double-blind, placebo-controlled trials supporting the therapeutic effect of daily SSRIs on PE (26). Open-design studies and studies using subjective reporting or questionnaires showed greater variation in ejaculation delay than double-blind studies in which the ejaculation delay was prospectively assessed with a stopwatch.

Based on this meta-analysis, SSRIs were expected to increase the geometric mean IELT by 2.6-fold to 13.2-fold. Paroxetine was found to be superior to fluoxetine, clomipramine and sertraline. Sertraline was superior to fluoxetine, whereas the efficacy of clomipramine was not significantly different from fluoxetine and sertraline. Paroxetine was evaluated in doses of 20-40 mg, sertraline 25-200 mg, fluoxetine 10-60 mg and clomipramine 25-50 mg; there was no significant relationship between dose and response among the various drugs. There is limited evidence that citalopram may be less efficacious compared to other SSRIs, while fluvoxamine may not be effective (27,28).

Ejaculation delay may start a few days after drug intake, but it is more evident after 1 to 2 weeks since receptor desensitisation requires time to occur. While efficacy may be maintained for several years, tachyphylaxis (decreasing response to a drug following chronic administration) may occur after 6 to 12 months (24).

Common side-effects of SSRIs include fatigue, drowsiness, yawning, nausea, vomiting, dry mouth, diarrhoea and perspiration; they are usually mild and gradually improve after 2 to 3 weeks (24). Decreased libido, anorgasmia, anejaculation and ED have been also reported.

In one controlled trial, on-demand use of clomipramine (but not paroxetine), 3 to 5 h before intercourse, was reported to be efficacious, though IELT improvement was inferior compared to daily treatment with the same drug (29). However, on-demand treatment may be combined with an initial trial of daily treatment or concomitant low-dose daily treatment reducing adverse effects (30,31).

#### 4.8.3.1 *Dapoxetine*

Dapoxetine is a potent SSRI, which has been designed as an on-demand oral treatment for PE. It is quickly absorbed with a T<sub>max</sub> of 1.5 h and is rapidly cleared, avoiding accumulation.

An integrated analysis of two, double-blind, randomised, controlled trials (1,958 patients) with dapoxetine was published (32). Dapoxetine, 30 and 60 mg, was administered 1 to 3 h before intercourse. Intravaginal ejaculatory latency time improved from a baseline of 0.9 min to 1.75 min, 2.78 min and 3.32 min in the patient groups treated with placebo, 30 mg dapoxetine, and 60 mg dapoxetine, respectively. Improved ejaculation control was reported by 51% and 58% of patients in the 30 mg and 60 mg groups, respectively. Both dapoxetine doses were effective on the first dose. Common adverse events for 30 mg and 60 mg doses of dapoxetine, respectively, were nausea (8.7%, 20.1%), diarrhoea (3.9%, 6.8%), headache (5.9%, 6.8%), and dizziness (3.0%, 6.2%).

In a subanalysis of these two studies (33), 32% of men reported a two-category (from a 5-point scale, 'very poor' to 'very good') or greater increase in control and satisfaction with sexual intercourse after treatment. More than 95% of those men rated their PE as 'slightly better', 'better', or 'much better' on the global impression of change (7-point scale, 'much worse' to 'much better') while 67.1% gave ratings of 'better' or 'much better.' They also had greater improvements in IELT than men with less than a two-category increase in control, with a mean (SD) change from baseline of 3.7 (4.3) vs 0.77 (1.8) min, respectively. The proportions of men with a two-category or greater increase in control with dapoxetine 30 and 60 mg were 36.3% and 44.5%, respectively (vs 15% with placebo).

In another randomised, double-blind, parallel-group, placebo-controlled, phase II trial including 1,162 men in 22 countries (34), mean average IELT increased from 0.9 min at baseline (all groups) to 1.9 min, 3.2 min, and 3.5 min with placebo and dapoxetine 30 mg and dapoxetine 60 mg, respectively, at study end point. The geometric mean IELT increased from 0.7 min at baseline to 1.1 min, 1.8 min, and 2.3 min, respectively, at study end point. All PEP measures and IELTs improved significantly with dapoxetine versus placebo at week 12 and week 24 ( $p < 0.001$  for all). The most common adverse effects were nausea, dizziness, diarrhea, and headache. Adverse effects led to discontinuation in 1.3%, 3.9%, and 8.2% of subjects with placebo and dapoxetine 30 mg. Finally, in a randomised, double-blind, placebo controlled, phase III trial (1,238 men in USA and Canada), dapoxetine reduced the personal distress and interpersonal difficulty associated with PE (35).

Dapoxetine has been approved (December 2008) for the on-demand treatment of PE in seven European countries (Sweden, Austria, Finland, Germany, Spain, Italy and Portugal). This is currently the first and only drug approved for such an indication.

#### 4.8.3.2 Recommendation on the treatment of PE

	LE	GR
Selective serotonin receptor inhibitors (SSRIs).	1A	A

#### 4.8.4 Phosphodiesterase type 5 inhibitors

Several recent studies have supported the therapeutic role of PDE5 inhibitors in PE. They may reduce performance anxiety due to better erections and may down-regulate the erectile threshold to a lower level of arousal so that greater arousal is required to achieve the ejaculation threshold. However, many of the mechanisms involved remain speculative (33,36-38).

There is only one well-designed, randomised, double-blind, placebo-controlled study comparing sildenafil to placebo (39). Although IELT was not significantly improved, sildenafil increased confidence, the perception of ejaculatory control and overall sexual satisfaction, reduced anxiety and decreased the refractory time to achieve a second erection after ejaculation.

In another randomised, double-blind, placebo-controlled study, lidocaine-prilocaine had similar efficacy to combination with sildenafil (50 mg before intercourse), while the efficacy of sildenafil was similar to placebo (no IELT data provided) (16). In contrast, in a randomised, double-blind, parallel group study, sildenafil significantly improved IELT and satisfaction and reduced overall anxiety compared to several SSRIs and the 'pause-squeeze' technique. From a baseline of IELT at 1 min, IELT improved to 15 min with sildenafil, 4 min with clomipramine, 3 min with sertraline, 4 min with paroxetine and 3 min with the 'pause-squeeze' technique (5).

Finally, several open-label studies showed that sildenafil combined with an SSRI is superior to SSRI monotherapy. Sildenafil combined with paroxetine improved IELT significantly and satisfaction versus paroxetine alone (40). Sildenafil combined with sertraline improved IELT and satisfaction significantly versus sertraline alone (41). Sildenafil combined with paroxetine and psychological and behavioural counselling significantly improved IELT and satisfaction in patients in whom other treatments failed (42). Finally, sildenafil combined with behavioural therapy significantly improved IELT and satisfaction versus behavioural therapy alone (43).

There are limited data on the efficacy in PE of other PDE5 inhibitors (tadalafil and vardenafil) (37, 38). Overall, the role of PDE5 inhibitors in PE patients without ED is not established, with only minimal double-blind placebo controlled data are available.

#### 4.8.4.1 Recommendation on the use of PDE5 inhibitors for the treatment of PE

	LE	GR
In patients presenting with PE, a trial of PDE5 inhibitors may be attempted.	2B	C

#### 4.8.5 Other drugs

Adrenergic blockade for PE aims to decrease the sympathetic tone of the seminal tract and therefore delay ejaculation (44). Tramadol is a centrally acting analgesic agent that combines opioid receptor activation and re-uptake inhibition of serotonin and noradrenaline.

Research suggests that the alpha-1 adrenergic antagonists, terazosin and alfuzosin (45,46), and tramadol (47,48) may have some efficacy in PE. However, further research is needed to investigate their role fully. Currently they are not recommended in clinical practice (49).


#### 4.8.6 Recommendations on the treatment of PE

	LE	GR
ED, other sexual dysfunction or genitourinary infection (e.g. prostatitis) should be treated first.	2a	B
Behavioural techniques have demonstrated benefit in treating PE. However, they are time intensive, require the support of a partner and can be difficult to do.	3	C
Pharmacotherapy is the basis of treatment in lifelong PE.	1a	A

Daily SSRIs are first-line, off-label, pharmacological treatment for PE. The pharmacokinetic profile of SSRIs is not amenable to pm dosing.	1a	A
Dapoxetine, a short-acting SSRI, has already been approved for the on-demand treatment of PE in seven European Countries.	1a	A
Topical anaesthetic agents provide viable alternatives to SSRIs (off-label).	1b	A
Recurrence is likely after treatment cessation.	1b	A
Behavioural therapy may augment pharmacotherapy to enhance relapse prevention.	3	C

*ED = erectile dysfunction; PE = premature ejaculation; SSRI = selective serotonin reuptake inhibitor; pm = on-demand administration.*

**Figure 4: Management of PE\***



\* Adapted from Lue et al. 2004 (49).

ED = erectile dysfunction; PE = premature ejaculation; IELT = intravaginal ejaculatory latency time; SSRI = selective serotonin receptor inhibitor.



#### 4.9 References

1. Semans JH. Premature ejaculation: a new approach. *South Med J* 1956 Apr;49(4):353-8. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/13311629>
2. de Carufel F, Trudel G. Effects of a new functional-sexological treatment for premature ejaculation. *J Sex Marital Ther* 2006 Mar-Apr;32(2):97-114.  
<http://www.ncbi.nlm.nih.gov/pubmed/16418103>
3. Grenier G, Byers ES. Rapid ejaculation: a review of conceptual, etiological, and treatment issues. *Arch Sex Behav* 1995 Aug;24(4):447-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/7661658>
4. Metz ME, Pryor JL, Nesvaci LJ, et al. Premature ejaculation: a psychophysiological review. *J Sex Marital Ther* 1997 Spring;23(1):3-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/9094032>
5. Abdel-Hamid IA, El Naggar EA, El Gilany AH. Assessment of as needed use of pharmacotherapy and the pause-squeeze technique in premature ejaculation. *Int J Impot Res* 2001 Feb;13(1):41-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/11313839>
6. De Amicis LA, Goldberg DC, LoPiccolo J, et al. Clinical follow-up of couples treated for sexual dysfunction. *Arch Sex Behav* 1985 Dec;14(6):467-89.  
<http://www.ncbi.nlm.nih.gov/pubmed/4084048>
7. Hawton K, Catalan J, Martin P, et al. Long-term outcome of sex therapy. *Behav Res Ther* 1986; 24(6):665-75. [no abstract available]  
<http://www.ncbi.nlm.nih.gov/pubmed/3800838>
8. Morales A, Barada J, Wyllie MG. A review of the current status of topical treatments for premature ejaculation. *BJU Int* 2007 Sep;100(3):493-501.  
<http://www.ncbi.nlm.nih.gov/pubmed/17608824>
9. Sachs BD, Liu YC. Maintenance of erection of penile glans, but not penile body, after transection of rat cavernous nerves. *J Urol* 1991 Sep;146(3):900-5.  
<http://www.ncbi.nlm.nih.gov/pubmed/1875517>
10. Wieder JA, Brackett NL, Lynne CM, et al. Anesthetic block of the dorsal penile nerve inhibits vibratory-induced ejaculation in men with spinal cord injuries. *Urology* 2000 Jun;55(6):915-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10840108>
11. Atikeler MK, Gecit I, Senol FA. Optimum usage of prilocaine-lidocaine cream in premature ejaculation. *Andrologia*. 2002 Dec;34(6):356-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12472618>
12. Busato W, Galindo CC. Topical anaesthetic use for treating premature ejaculation: a double-blind, randomized, placebo-controlled study. *BJU Int* 2004 May;93(7):1018-21.  
<http://www.ncbi.nlm.nih.gov/pubmed/15142155>
13. Henry R, Morales A, Wyllie MG. TEMPE: Topical Eutectic-Like Mixture for Premature Ejaculation. *Expert Opin Drug Deliv* 2008 Feb;5(2):251-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/18248322>
14. Dinsmore WW, Hackett G, Goldmeier D, et al. Topical eutectic mixture for premature ejaculation (TEMPE): a novel aerosol-delivery form of lidocaine-prilocaine for treating premature ejaculation. *BJU Int* 2007 Feb;99(2):369-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/17129234>
15. Henry R, Morales A. Topical lidocaine-prilocaine spray for the treatment of premature ejaculation: a proof of concept study. *Int J Impot Res* 2003 Aug;15(4):277-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/12934056>
16. Atan A, Basar MM, Tuncel A, et al. Comparison of efficacy of sildenafil-only, sildenafil plus topical EMLA cream, and topical EMLA-cream-only in treatment of premature ejaculation. *Urology* 2006 Feb; 67(2):388-91.  
<http://www.ncbi.nlm.nih.gov/pubmed/16461091>
17. Xin ZC, Choi YD, Seong DH, et al. Sensory evoked potential and effect of SS-cream in premature ejaculation. *Yonsei Med J* 1995 Nov;36(5):397-401.  
<http://www.ncbi.nlm.nih.gov/pubmed/8545998>
18. Xin ZC, Choi YD, Lee WH, et al. Penile vibratory threshold changes with various doses of SS-cream in patients with primary premature ejaculation. *Yonsei Med J* 2000 Feb;41(1):29-33.  
<http://www.ncbi.nlm.nih.gov/pubmed/10731916>
19. Choi HK, Jung GW, Moon KH, et al. Clinical study of SS-cream in patients with lifelong premature ejaculation. *Urology* 2000 Feb;55(2):257-61.  
<http://www.ncbi.nlm.nih.gov/pubmed/10688090>

20. Truitt WA, Coolen LM. Identification of a potential ejaculation generator in the spinal cord. *Science* 2002 Aug 30;297(5586):1566-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12202834>
21. Borgdorff AJ, Bernabé J, Denys P, et al. Ejaculation elicited by microstimulation of lumbar spinothalamic neurons. *Eur Urol* 2008 Aug;54(2):449-56.  
<http://www.ncbi.nlm.nih.gov/pubmed/18394782>
22. Giuliano F. 5-Hydroxytryptamine in premature ejaculation: opportunities for therapeutic intervention. *Trends Neurosci* 2007 Feb;30(2):79-84.  
<http://www.ncbi.nlm.nih.gov/pubmed/17169440>
23. Olivier B, van Oorschoot R, Waldinger MD. Serotonin, serotonergic receptors, selective serotonin reuptake inhibitors and sexual behaviour. *Int Clin Psychopharmacol* 1998 Jul;13 Suppl 6:S9-14.  
<http://www.ncbi.nlm.nih.gov/pubmed/9728669>
24. Waldinger MD. Premature ejaculation: definition and drug treatment. *Drugs* 2007;67(4):547-68.  
<http://www.ncbi.nlm.nih.gov/pubmed/17352514>
25. Waldinger MD, Hengeveld MW, Zwinderman AH. Paroxetine treatment of premature ejaculation: a double-blind, randomized, placebo-controlled study. *Am J Psychiatry* 1994 Sep;151(1):1377-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/8067497>
26. Waldinger MD, Zwinderman AH, Schweitzer DH, et al. Relevance of methodological design for the interpretation of efficacy of drug treatment of premature ejaculation: a systematic review and meta-analysis. *Int J Impot Res* 2004 Aug;16(4):369-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/14961051>
27. Waldinger MD, Zwinderman AH, Olivier B. SSRIs and ejaculation: a double-blind, randomized, fixed-dose study with paroxetine and citalopram. *J Clin Psychopharmacol* 2001 Dec;21(6):556-60.  
<http://www.ncbi.nlm.nih.gov/pubmed/11763001>
28. Waldinger MD, Hengeveld MW, Zwinderman AH, et al. Effect of SSRI antidepressants on ejaculation: a double-blind, randomized, placebo-controlled study with fluoxetine, fluvoxamine, paroxetine, and sertraline. *J Clin Psychopharmacol* 1998 Aug;18(4):274-81.  
<http://www.ncbi.nlm.nih.gov/pubmed/9690692>
29. Waldinger MD, Zwinderman AH, Olivier B. On-demand treatment of premature ejaculation with clomipramine and paroxetine: a randomized, double-blind fixed-dose study with stopwatch assessment. *Eur Urol* 2004 Oct;46(4):510-5; discussion 516.  
<http://www.ncbi.nlm.nih.gov/pubmed/15363569>
30. McMahon CG, Touma K. Treatment of premature ejaculation with paroxetine hydrochloride as needed: 2 single-blind placebo controlled crossover studies. *J Urol* 1999 Jun;161(6):1826-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/10332446>
31. Kim SW, Paick JS. Short-term analysis of the effects of as needed use of sertraline at 5 PM for the treatment of premature ejaculation. *Urology* 1999 Sep;54(3):544-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/10475369>
32. Pryor JL, Althof SE, Steidle C, et al; Dapoxetine Study Group. Efficacy and tolerability of dapoxetine in treatment of premature ejaculation: an integrated analysis of two double-blind, randomised controlled trials. *Lancet* 2006 Sep;368(9539):929-37.  
<http://www.ncbi.nlm.nih.gov/pubmed/16962882>
33. Shabsigh R, Patrick DL, Rowland DL, et al. Perceived control over ejaculation is central to treatment benefit in men with premature ejaculation: results from phase III trials with dapoxetine. *BJU Int* 2008 Sep;102(7):824-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/18647300>
34. Buvat J, Tesfaye F, Rothman M, et al. Dapoxetine for the treatment of premature ejaculation: results from a randomized, double-blind, placebo-controlled phase 3 trial in 22 countries. *Eur Urol* 2009 Apr;55(4):957-67.  
<http://www.ncbi.nlm.nih.gov/pubmed/19195772>
35. Kaufman JM, Rosen RC, Mudumbi RV, et al. Treatment benefit of dapoxetine for premature ejaculation: results from a placebo-controlled phase III trial. *BJU Int* 2009 Mar;103(5):651-8.  
<http://www.ncbi.nlm.nih.gov/pubmed/19021601>
36. Chen J, Keren-Paz G, Bar-Yosef Y, et al. The role of phosphodiesterase type 5 inhibitors in the management of premature ejaculation: a critical analysis of basic science and clinical data. *Eur Urol* 2007 Nov;52(5):1331-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/17728050>
37. McMahon CG, McMahon CN, Leow LJ, et al. Efficacy of type-5 phosphodiesterase inhibitors in the drug treatment of premature ejaculation: a systematic review. *BJU Int* 2006 Aug;98(2):259-72.  
<http://www.ncbi.nlm.nih.gov/pubmed/16879663>

38. Wang WF, Minhas S, Ralph DJ. Phosphodiesterase 5 inhibitors in the treatment of premature ejaculation. *Int J Androl* 2006 Oct;29(5):503-09.  
<http://www.ncbi.nlm.nih.gov/pubmed/16573707>
39. McMahon CG, Stuckey BG, Andersen M, et al. Efficacy of sildenafil citrate (Viagra) in men with premature ejaculation. *J Sex Med* 2005 May;2(3):368-75.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422868>
40. Salonia A, Maga T, Colombo R, et al. A prospective study comparing paroxetine alone versus paroxetine plus sildenafil in patients with premature ejaculation. *J Urol* 2002 Dec;168(6):2486-9.  
<http://www.ncbi.nlm.nih.gov/pubmed/12441946>
41. Zhang XS, Wang YX, Huang XY, et al. [Comparison between sildenafil plus sertraline and sertraline alone in the treatment of premature ejaculation]. *Zhonghua Nan Ke Xue* 2005 Jul;11(7):520-2,525. [Article in Chinese]  
<http://www.ncbi.nlm.nih.gov/pubmed/16078671>
42. Chen J, Mabjeesh NJ, Matzkin H, et al. Efficacy of sildenafil as adjuvant therapy to selective serotonin reuptake inhibitor in alleviating premature ejaculation. *Urology* 2003 Jan;61(1):197-200.  
<http://www.ncbi.nlm.nih.gov/pubmed/12559295>
43. Tang W, Ma L, Zhao L, et al. [Clinical efficacy of Viagra with behavior therapy against premature ejaculation]. *Zhonghua Nan Ke Xue* 2004 May;10(5):366-7, 370. [Article in Chinese]  
<http://www.ncbi.nlm.nih.gov/pubmed/15190831>
44. Hsieh JT, Chang HC, Law HS, et al. In vivo evaluation of serotonergic agents and alpha-adrenergic blockers on premature ejaculation by inhibiting the seminal vesicle pressure response to electrical nerve stimulation. *Br J Urol* 1998 Aug;82(2):237-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/9722759>
45. Başar MM, Yilmaz E, Ferhat M, et al. Terazosin in the treatment of premature ejaculation: a short-term follow-up. *Int Urol Nephrol* 2005;37(4):773-7.  
<http://www.ncbi.nlm.nih.gov/pubmed/16362597>
46. Cavallini G. Alpha-1 blockade pharmacotherapy in primitive psychogenic premature ejaculation resistant to psychotherapy. *Eur Urol* 1995;28(2):126-30.  
<http://www.ncbi.nlm.nih.gov/pubmed/8529737>
47. Safarinejad MR, Hosseini SY. Safety and efficacy of tramadol in the treatment of premature ejaculation: a double-blind, placebo-controlled, fixed-dose, randomized study. *J Clin Psychopharmacol* 2006 Feb;26(1):27-31.  
<http://www.ncbi.nlm.nih.gov/pubmed/16415702>
48. Salem EA, Wilson SK, Bissada NK, et al. Tramadol HCL has promise in on-demand use to treat premature ejaculation. *J Sex Med* 2008 Jan;5(1):188-93.  
<http://www.ncbi.nlm.nih.gov/pubmed/17362279>
49. Lue TF, Giuliano F, Montorsi F, et al. Summary of the recommendations on sexual dysfunctions in men. *J Sex Med* 2004 Jul;1(1):6-23.  
<http://www.ncbi.nlm.nih.gov/pubmed/16422979>

## 5. CONCLUSION

Modern treatment of ED has been revolutionised by the worldwide availability of three PDE5Is for oral use – sildenafil, tadalafil and vardenafil. These drugs have high efficacy and safety rates, even in difficult-to-treat populations, such as patients with diabetes mellitus or who have undergone RP. Patients should be encouraged to try all three PDE5Is. Patients should make up their own minds about which compound has the best efficacy, while also considering other factors, such as time of onset, duration of action, window of opportunity and how side-effects affect them individually.

Treatment options for patients who do not respond to oral drugs, or for whom drugs are contraindicated, include intracavernous injections, intraurethral alprostadil, vacuum constriction devices, or implantation of a penile prosthesis.

It is very important that the physician warns the patient that sexual intercourse is a vigorous physical activity, which increases heart rate as well as cardiac work. Physicians should assess the cardiac fitness of patients prior to treating ED.

Any successful pharmacological treatment for erectile failure demands a degree of integrity of the penile mechanisms of erection. Further studies of individual agents and synergistic activity of available substances are underway. The search for the ideal pharmacological therapy for erectile failure aims to fulfil the following characteristics: good efficacy, easy administration, freedom from toxicity and side-effects, with a rapid onset and a possible long-acting effect.

Premature ejaculation is another very common male sexual dysfunction, with prevalence rates of 20% to 30%. Four major definitions of PE are currently used and the most widely accepted classification of PE includes “lifelong” (primary) and “acquired” (secondary) forms (syndromes).

Diagnosis of PE in everyday clinical practice is based on medical and sexual history assessing IELT, perceived control, distress, and interpersonal difficulty related to the ejaculatory dysfunction. Physical examination and laboratory testing may be needed in selected patients only.

Pharmacotherapy is the basis of treatment in lifelong PE including daily dosing of SSRIs and topical anaesthetics. Behavioural techniques may be efficacious as a monotherapy or in combination with pharmacotherapy, but they can be difficult to perform. In every case, recurrence is likely to occur after treatment withdrawal.

## 6. ABBREVIATIONS USED IN THE TEXT

*This list is not comprehensive for the most common abbreviations*

5-HT	5-hydroxytryptamine
AIPE	Arabic Index of Premature Ejaculation
AUC	area under curve - serum concentration time curve
BMI	body mass index
CAD	coronary artery disease
cGMP	cyclic guanosine monophosphate
CGRP	calcitonin gene-related peptide
CHF	congestive heart failure
C <sub>max</sub>	maximal concentration
DICC	dynamic infusion cavernosometry or cavernosography
DRE	digital rectal examination
DSM-IV-TR	Diagnostic and Statistical Manual of Mental Disorders IV-Text Revision
EAU	European Association of Urology
ED	erectile dysfunction
EMA	European Medicines Agency
FDA	(US) Food and Drug Administration
FSH	follicle-stimulating hormone
GAQ	General Assessment Question
GR	grade of recommendation
GSSAB	Global Study of Sexual Attitudes and Behaviors
ICD-10	International Classification of Diseases-10
IELT	intravaginal ejaculatory latency time
IIEF	International Index for Erectile Function
IIEF-EF	International Index for Erectile Function - erectile function domain
IPE	Index of Premature Ejaculation
ISSM	International Society for Sexual Medicine
LE	level of evidence
LH	luteinising hormone
LVD	left ventricular dysfunction
MET	metabolic equivalent of energy expenditure in the resting state
MI	myocardial infarction
MMAS	Massachusetts Male Aging Study
MSHQ-EjD	Male Sexual Health Questionnaire Ejaculatory Dysfunction
NHLS	National Health and Social Life Survey
NS	nerve sparing
NO	nitric oxide
NPTR	nocturnal penile tumescence and rigidity
NSRP	nerve-sparing radical prostatectomy
NYHA	New York Heart Association
PCa	prostate cancer
PDE5Is	phosphodiesterase type 5 inhibitors
PE	premature ejaculation
PEDT	Premature Ejaculation Diagnostic Tool
PEP	Premature Ejaculation Profile
PEPA	Premature Ejaculation Prevalence and Attitudes
PRO	Patient reported outcome
PSA	prostate-specific antigen
QoL	quality of life
RP	radical prostatectomy
SEP	sexual encounter profile
SSRI	selective serotonin reuptake inhibitor
TEMPE	topical eutectic mixture for premature ejaculation
T <sub>max</sub>	time to maximum plasma concentration
VCD	vacuum constriction devices
VIP	vasointestinal peptide

### **Conflict of interest**

All members of the Male Sexual Dysfunction Guidelines working panel have provided disclosure statements on all relationships that they have that might be perceived to be a potential source of a conflict of interest. This information is publically accessible through the European Association of Urology website. This guidelines document was developed with the financial support of the European Association of Urology. No external sources of funding and support have been involved. The EAU is a non-profit organisation, and funding is limited to administrative assistance and travel and meeting expenses. No honoraria or other reimbursements have been provided.